DUTCH BIRDING

VOLUME 28 • NO 6 • 2006

Dutch Birding

Internationaal tijdschrift over Palearctische vogels

Redactie
Dutch Birding
Postbus 116
2080 AC Santpoort-Zuid
Nederland
fax 023-5376749
e-mail editors@dutchbirding.nl

Fotoredactie
Dutch Birding
p/a René Pop
Postbus 31
1790 AA Den Burg-Texel
Nederland
e-mail rene.pop@dutchbirding.nl

Abonnementenadministratie
p/a Jeannette Admiraal
Iepenlaan 11
1901 ST Castricum
Nederland
e-mail circulation@dutchbirding.nl

Bestuur
Dutch Birding Association
Postbus 75611
1070 AP Amsterdam
Nederland
e-mail dba@dutchbirding.nl

Commissie Dwaalgasten Nederlandse Avifauna
CDNA
Postbus 45
2080 AA Santpoort-Zuid
Nederland
e-mail cdna@dutchbirding.nl

Commissie Systematiek Nederlandse Avifauna
CSNA, p/a George Sangster
Stevenshof 17
2312 GM Leiden
Nederland
e-mail csna@dutchbirding.nl

Telefoonlijnen
0900-BIRDING (= 0900-2473464)
(vogellijn, EUR 0.35/min)
010-4281212 (inspreeklijn)

Internet
www.dutchbirding.nl

Dutch Birding

Hoofdredacteur Arnoud van den Berg (tel 023-5378024, e-mail arnoud.van.den.berg@dutchbirding.nl)

Adjunct hoofdredacteur Enno Ebels (tel 030-2961335, e-mail enno.ebels@dutchbirding.nl)

Uitvoerend redacteur André van Loon (tel / fax 020-6997585, e-mail andre.van.loon@dutchbirding.nl)

Fotografisch redacteur René Pop (tel 0222-316801, fax 0222-316802, e-mail rene.pop@dutchbirding.nl)

Redactieraad Peter Adriaens, Ferdy Hieselaar, Roy Slaterus, Roland van der Vliet en Rik Winters

Redactie-adviesraad Peter Barthel, Mark Constantine, Gunter De Smet, Dick Forsman, Ricard Gutiérrez, Anthony McGeehan, Killian Mullarney, Klaus Malling Olsen, Magnus Robb, George Sangster, Hadoram Shirihai, Brian Small en Lars Svensson

Redactiemedewerkers Rob van Bemmelen, Marten van Dijl, Ruud van Dongen, Gerald Driessens, Nils van Duivendijk, Joris Elst, Dick Groenendijk, Marcel Haas, Jan van der Laan, Hans van der Meulen, André van der Plas, Kees Roselaar, René van Rossum en Peter de Rouw

Productie en lay-out André van Loon en René Pop

Advertenties Laurens Steijn, p/a Dutch Birding, Postbus 75611, 1070 AP Amsterdam e-mail advertising@dutchbirding.nl

Abonnementen De abonnementsprijs voor 2006 bedraagt: EUR 32.00 (Nederland), EUR 35.00 (Europa) en EUR 38.00 (landen buiten Europa).
U kunt zich abonneren door het overmaken van de abonnementsprijs op girorekening 01 50 697 (Nederland), girorekening 000 1592468 19 (België) of bankrekening 54 93 30 348 van ABN•AMRO (Castricum), ovv 'abonnement Dutch Birding'. Alle rekeningen zijn ten name van de Dutch Birding Association. Het abonnement gaat in na ontvangst van de betaling.

Dutch Birding is een tweemaandelijks tijdschrift. Het publiceert originele artikelen en mededelingen over morfologie, systematiek, voorkomen en verspreiding van vogels in de Benelux, Europa en elders in het Palearctische gebied. Het publiceert tevens bijdragen over vogels in het Aziatisch-Pacifische gebied en andere gebieden.

De volgorde van vogels in Dutch Birding volgt in eerste instantie een klassieke 'Wetmore-indeling'. Binnen dit raamwerk worden voor taxonomie en naamgeving de volgende overzichten aangehouden: *Dutch Birding-namen* door A B van den Berg (2006, Amsterdam) (taxonomie en wetenschappelijke, Nederlandse en Engelse namen van West-Palearctische vogels); *Vogels van de wereld – complete checklist* door M Walters (1997, Baarn) (Nederlandse namen van overige vogels van de wereld); en *The Howard and Moore complete checklist of the birds of the world* (derde editie) door E C Dickinson (redactie) (2003, Londen) (taxonomie en wetenschappelijke en Engelse namen van overige vogels van de wereld).

Een lijst met tarieven voor de vergoeding van auteurs, fotografen en tekenaars is verkrijgbaar bij de redactie.

Dutch Birding Association

Bestuur Theo Admiraal (penningmeester), Gijsbert van der Bent (voorzitter, tel 071-4024547), Arjan van Egmond, Wietze Janse, Marc Plomp (secretaris, tel 0348-433730) en Laurens Steijn; tevens is de redactie van Dutch Birding met een zetel vertegenwoordigd.

Bestuursmedewerkers Jeannette Admiraal, Leon Boon, Albert van den Ende, Leo Heemskerk, Arnold Meijer, Chris van Rijswijk, Henk van Rijswijk, Willem van Rijswijk, Vincent van der Spek, Michel Veldt, Jeroen van Vianen en Peter Weiland.

Commissie Dwaalgasten Nederlandse Avifauna (CDNA)

Leden Ruud Brouwer, Nils van Duivendijk, Dick Groenendijk, Teus Luijendijk, Laurens Steijn (secretaris, tel 020-7765127), Roland van der Vliet (voorzitter, tel 030-2970013), Arend Wassink en Pim Wolf. **Medewerker** Max Berlijn (archivaris). De CDNA is een commissie van de Dutch Birding Association en de Nederlandse Ornithologische Unie.

Commissie Systematiek Nederlandse Avifauna (CSNA)

Leden Arnoud van den Berg, André van Loon, Kees Roselaar en George Sangster (secretaris, tel 071-5143790). De CSNA is een commissie van de Dutch Birding Association en de Nederlandse Ornithologische Unie.

© 2006 Stichting Dutch Birding Association. Het copyright van de foto's en tekeningen blijft bij de fotografen en tekenaars. ISSN 0167-2878.

Drukkerij robstolk®, Mauritskade 55, 1092 AD Amsterdam, Nederland

Dutch Birding

Chief editor Arnoud van den Berg (tel +31-235378024, e-mail arnoud.van.den.berg@dutchbirding.nl)

Deputy chief editor Enno Ebels (tel +31-302961335, e-mail enno.ebels@dutchbirding.nl)

Executive editor André van Loon (tel / fax +31-206997585, e-mail andre.van.loon@dutchbirding.nl)

Photographic editor René Pop (tel +31-222316801, fax +31-222316802, e-mail rene.pop@dutchbirding.nl)

Editorial board Peter Adriaens, Ferdy Hieselaar, Roy Slaterus, Roland van der Vliet and Rik Winters

Editorial advisory board Peter Barthel, Mark Constantine, Gunter De Smet, Dick Forsman, Ricard Gutiérrez, Anthony McGeehan, Killian Mullarney, Klaus Malling Olsen, Magnus Robb, George Sangster, Hadoram Shirihai, Brian Small and Lars Svensson

Editorial assistants Rob van Bemmelen, Marten van Dijl, Ruud van Dongen, Gerald Driessens, Nils van Duivendijk, Joris Elst, Dick Groenendijk, Marcel Haas, Jan van der Laan, Hans van der Meulen, André van der Plas, Kees Roselaar, René van Rossum and Peter de Rouw

Production and lay-out André van Loon and René Pop

Advertising Laurens Steijn, c/o Dutch Birding, Postbus 75611, 1070 AP Amsterdam e-mail advertising@dutchbirding.nl

Subscriptions The subscription rate for 2006 is: EUR 32.00 (Netherlands), EUR 35.00 (Europe) and EUR 38.00 (countries outside Europe).
Subscribers in Belgium, Denmark, Finland, Germany, Norway and Sweden are kindly requested to pay the subscription fee to our local bank accounts in these countries. Details can be found on the internet (www.dutchbirding.nl).
British subscribers are requested to pay exclusively by Sterling cheque. This cheque can be sent to Dutch Birding, c/o Jeannette Admiraal, Iepenlaan 11, 1901 ST Castricum, Netherlands.
Subscribers in other countries can make their payment by credit card (Access, Eurocard, MasterCard or Visa). Please send an e-mail indicating your credit card type, account number, the expiry date and full address details to circulation@dutchbirding.nl.
The subscription starts upon receipt of payment and already published issues will be sent.

Dutch Birding is a bimonthly journal. It publishes original papers and notes on morphology, systematics, occurrence and distribution of birds in the Benelux, Europe and elsewhere in the Palearctic region. It also publishes contributions on birds in the Asian-Pacific region and other regions.

The sequence of birds in Dutch Birding basically follows a classic 'Wetmore sequence'. Within this framework, the following lists are used for taxonomy and nomenclature: *Dutch Birding's names* by A B van den Berg (2006, Amsterdam) (taxonomy and scientific, Dutch and English names of Western Palearctic birds); *Vogels van de wereld – complete checklist* by M Walters (1997, Baarn) (Dutch names of remaining birds of the world); and *The Howard and Moore complete checklist of the birds of the world* (third edition) by E C Dickinson (editor) (2003, London) (taxonomy and scientific and English names of remaining birds of the world).

A schedule of payment rates for authors, photographers and artists is available from the editors.

Dutch Birding Association

Board Theo Admiraal (treasurer), Gijsbert van der Bent (president, tel +31-714024547), Arjan van Egmond, Wietze Janse, Marc Plomp (secretary, tel +31-348433730) and Laurens Steijn; the editors of Dutch Birding also have one seat in the board.

Board assistants Jeannette Admiraal, Leon Boon, Albert van den Ende, Leo Heemskerk, Arnold Meijer, Chris van Rijswijk, Henk van Rijswijk, Willem van Rijswijk, Vincent van der Spek, Michel Veldt, Jeroen van Vianen and Peter Weiland.

Dutch rarities committee (CDNA)

Members Ruud Brouwer, Nils van Duivendijk, Dick Groenendijk, Teus Luijendijk, Laurens Steijn (secretary, tel +31-207765127), Roland van der Vliet (chairman, tel +31-302970013), Arend Wassink and Pim Wolf. **Assistant** Max Berlijn (archivist). The CDNA is a committee of the Dutch Birding Association and the Netherlands Ornithologists' Union.

Dutch committee for avian systematics (CSNA)

Members Arnoud van den Berg, André van Loon, Kees Roselaar and George Sangster (secretary, tel +31-715143790). The CSNA is a committee of the Dutch Birding Association and the Netherlands Ornithologists' Union.

© 2006 Stichting Dutch Birding Association. The copyright of the photographs and drawings remains with the photographers and artists. ISSN 0167-2878.

Printed by drukkerij robstolk®, Mauritskade 55, 1092 AD Amsterdam, Netherlands

Dutch Birding

International journal on Palearctic birds

Editors
Dutch Birding
Postbus 116
2080 AC Santpoort-Zuid
Netherlands
fax +31-235376749
e-mail editors@dutchbirding.nl

Photographic editor
Dutch Birding
c/o René Pop
Postbus 31
1790 AA Den Burg-Texel
Netherlands
e-mail rene.pop@dutchbirding.nl

Subscription administration
c/o Jeannette Admiraal
Iepenlaan 11
1901 ST Castricum
Netherlands
e-mail circulation@dutchbirding.nl

Board
Dutch Birding Association
Postbus 75611
1070 AP Amsterdam
Netherlands
e-mail dba@dutchbirding.nl

Dutch rarities committee
CDNA
Postbus 45
2080 AA Santpoort-Zuid
Netherlands
e-mail cdna@dutchbirding.nl

Dutch committee for avian systematics
CSNA, c/o George Sangster
Stevenshof 17
2312 GM Leiden
Netherlands
e-mail csna@dutchbirding.nl

Internet
www.dutchbirding.nl

Dutch Birding JAARGANG 28 NUMMER 6 2006 *VOLUME 28 NUMBER 6 2006*

Artikelen / papers
345 Rare birds in the Netherlands in 2005 *Roland E van der Vliet, Jan van der Laan, Max Berlijn & CDNA*

Masters of Mystery
366 Solutions of fifth round 2006: Mediterranean Gull and Bateleur; Sixth round 2006 *Rob S A van Bemmelen & Dick Groenendijk*

CDNA-mededelingen
368 CDNA-postadres gewijzigd

Aankondigingen & verzoeken / announcements & requests
368 Ornithologisch jaarverslag Texel 2005

Corrigendum
368

DBA-nieuws
369 Betaling van abonnementsgeld voor 2007; Payment of subscription for 2007; Uitverkochte nummers op DBA-website en invoering digitaal abonnement; Sold-out back issues on DBA-website and introduction of digital subscription; Geanimeerde Texelweek 2006; Dutch Birding-vogelweek op 13-18 oktober 2007

WP reports
371 late September to early November 2006 *Arnoud B van den Berg & Marcel Haas*

Recente meldingen / recent reports
390 Nederland: september-oktober 2006 *Ruud M van Dongen, Remco Hofland & Peter W W de Rouw*
403 België: september-oktober 2006 *Gerald Driessens*

DB Actueel
407 Afwijkende spreeuw op Texel houdt gemoederen bezig [ABERRANT STARLING]; Vale Gierzwaluw op Vlieland: langverwachte nieuwkomer voor Nederland [PALLID SWIFT]; Eindelijk Rotszwaluwen [EURASIAN CRAG MARTINS]; Huisgierzwaluw te IJmuiden [LITTLE SWIFT]

Voorplaat / front cover
Roodstuittapuit / Red-rumped Wheatear *Oenanthe moesta*, Timahdite, Midden-Atlas, Marokko, 13 februari 2006 *(Daniele Occhiato)*

Rare birds in the Netherlands in 2005

Roland E van der Vliet, Jan van der Laan, Max Berlijn & CDNA

This is the 26th annual report on rare birds in the Netherlands to be published in Dutch Birding. This report comprises records from 2005 as well as belated and reconsidered records, dating back to 1890, which have been evaluated by the Dutch rarities committee, the Commissie Dwaalgasten Nederlandse Avifauna (CDNA). Several 2005 records are still under review or awaiting submission and could therefore not be included. In addition, some records for 2004 and earlier years are still under consideration for various reasons.

Details included for each record are, if available: date(s); location and/or *municipality* (municipal divisions as on 1 January 2005); province; number of birds if more than one, plumage and sex; type of record if trapped, photographed, sound-recorded, videoed or found dead (and where specimen is stored); names of up to three observers involved in finding, identifying and recording, and relevant references in the literature, which normally include (inter)national but not regional journals, and published photographs. Records from 2004 lasting into 2005 that were already published in the 2004 report (van der Vliet et al 2005) are repeated here without references, for the sake of completeness. Observations not (yet) submitted mentioned in the 2004 report are not repeated in this report and the CDNA recommends consulting the 2004 report for these. Sequence of the records in the species accounts is from the current year backwards, with records within one year presented chronologically. A compilation of most of the birds videoed can be found in Boon & Slaterus (2006) and Plomp et al (2006). Numbers after each (sub)species' name refer to the total number of individuals ***1*** from 1 January 1800 to 31 December 1979, ***2*** since 1 January 1980 but excluding ***3*** the current year. Taxa marked with an asterisk * are new to the Dutch list.

The following CDNA members voted on some or all of the records in this report: Theo Bakker, Max Berlijn, Ruud E Brouwer, A (Bert) de Bruin (former secretary), Nils van Duivendijk, Dick Groenendijk, Teus J C Luijendijk, Laurens B Steijn (secretary), Roland E van der Vliet (chairman), Arend Wassink and Pim A Wolf. Records can be submitted through email (cdna@dutchbirding.nl) or via the CDNA site on the Dutch Birding website (www.dutchbirding.nl), using an electronic submission form. Records can also be sent by ordinary mail to CDNA, Postbus 116, 2080 AC Santpoort-Zuid, the Netherlands, preferably using standard forms which can be obtained free of charge from the CDNA. On the website, it is possible to see which records are currently under review, which were recently accepted and which were rejected.

Decisions regarding taxonomy have been delegated to the Dutch committee for avian systematics, the Commissie Systematiek Nederlandse Avifauna (CSNA), which on 1 January 2005 consisted of the following members: Arnoud B van den Berg, André J van Loon, C S (Kees) Roselaar and George Sangster (secretary).

In 2005, at least 357 species were recorded, of which at least two are new to the Dutch list: Eastern Imperial Eagle *Aquila heliaca* and Killdeer *Charadrius vociferus*. A record of Purple-backed Starling *Sturnus sturninus*, another potential new species, is still under review (two previous records, both in 1999, have been regarded as escapes). From previous years, other additions to the Dutch list are Marbled Duck *Marmaronetta*

angustirostris (2004), Western Orphean Warbler *Sylvia hortensis* and Hornemann's Redpoll *Carduelis hornemanni hornemanni* (2003), Eurasian Pygmy Owl *Glaudicium passerinum* (2002) and Oriental Pratincole *Glareola maldivarum* (1997). Highlights from 2005 are the second Trumpeter Finch *Bucanetes githagineus*, the third Northern Hawk Owl *Surnia ulula* and Calandra Lark *Melanocorypha calandra*, the fourth White-throated Robin *Irania gutturalis* and Isabelline Wheatear *Oenanthe isabellina*, and the fifth Red-necked Stint *Calidris ruficollis*. 2005 was a record year for Desert Wheatear *O deserti* with nine records. In contrast, there were no records of vagrants which occurred almost annually in recent years, like American Golden Plover *Pluvialis dominica*, Sociable Lapwing *Vanellus gregarius*, Roseate Tern *Sterna dougallii* or Citrine Wagtail *Motacilla citreola*.

Reviews in progress involve records of Whistling Swan *Cygnus columbianus* (regarding identification) and Yellow-headed Blackbird *Xanthocephalus xanthocephalus* (regarding identification as well as status).

Systematic list of records

Ross's Goose / Ross' Gans
Anser rossii 0,5,1
2-21 January, Bochtjesplaat and Anjumerkolken, *Dongeradeel*, Friesland, adult, white morph, photographed (R Cazemier, D Hoekstra, T Bakker et al); 2 February to 18 April, Oostvaardersplassen, *Almere* and *Lelystad*, Flevoland, and Polder Zeevang, *Zeevang*, and Oosterpoel, *Waterland*, Noord-Holland, adult, white morph, photographed (R A van Dijk, D Tanger et al; Dutch Birding 27: 148, plate 187, 2005); 15 March to 29 April, Scherpenissepolder, *Tholen*, Zeeland, and *Goedereede*, and Westplaat Buitengronden, Melissant, *Dirksland*, Zuid-Holland, two (on at least 21-22 March), adult, white morph, photographed (P A Wolf et al); 27 April, Jan Durkspolder, Oudega, *Smallingerland*, Friesland, adult, white morph, photographed (J Huizenga); 19-22 October, Anjumerkolken, *Dongeradeel*, Friesland, adult, white morph, photographed (R Cazemier, O Tol et al).
2004 16 October to 21 February 2005, Rammegors and Scherpenissepolder, *Tholen*, and Yerseke, *Reimerswaal*, Zeeland, and Ouddorp, *Goedereede*, and Slikken van Flakkee, *Dirksland*, Zuid-Holland, adult, white morph, photographed.

On 21-22 March, three birds were simultaneously present in the Netherlands, which means that a presumed new bird had arrived, adding one bird to the totals.

Canvasback / Grote Tafeleend
Aythya valisineria 0,1,0
14 October to 8 November, Castricum aan Zee, *Castricum*, Noord-Holland, adult male, photographed, videoed (J Vroege, H Levering, R E Brouwer et al; Dutch Birding 27: 431, plate 540, 2005).

This is considered to be the returning bird first seen in January-March 2003 (Reijnders & Ebels 2004, van der Vliet et al 2004). It was not present (or, at least, not observed) in the winters of 2003/04 and 2004/05.

Ring-necked Duck / Ringsnaveleend
Aythya collaris 5,18,3
22 October to 22 January 2006, Budel-Dorplein, *Cranendonck*, Noord-Brabant, two (second bird from 29 October), first-winter female, photographed, videoed (R G Bouwman, H Verdaat, F Neijts et al; Dutch Birding 27: 431, plate 539, 2005); 4-6 December, Genne, *Hasselt*, Overijssel, adult female, photographed (D van de Zee, H B Bouman, G Veurink et al; Dutch Birding 28: 57, plate 75, 2006).

Remarkably, all three birds this year were females. Only two previous records concern females, in April 1996 and November 2001.

Ring-necked x Tufted Duck / Ringsnaveleend x Kuifeend
Aythya collaris x fuligula 0,6,0
2003 26 December to 2 March 2004, Dijkwielen, *Wieringermeer*, Noord-Holland, adult male, photographed (J Buijsman, R E Brouwer, H Niesen et al).

King Eider / Koningseider
Somateria spectabilis 1,10,1
14 March, Camperduin, *Bergen*, Noord-Holland, adult male (G J de Haan, N F van der Ham).

Previous records were in May 1975, December 1981 to March 1982, April 1989, October 1989 to April 1991 (two), March 1993 to May 1995, October-November 1994, May-December 1999, August 2000 to July 2001 (two) and May-September 2003. Only three records refer to females: in April 1989, March 1993 to May 1995, and October-November 1994.

Surf Scoter / Brilzee-eend
Melanitta perspicillata 3,13,3
13-14 March, Buren, *Ameland*, Friesland, two, adult male, photographed (M S J Hoekstein, P A Wolf, H Verdaat et al; Dutch Birding 27: 222, plate 271, 2005); 6-11 April, Brouwersdam, *Schouwen-Duiveland*, Zeeland, adult male, photographed (P A Wolf, P L Meininger, S J Lilipaly et al; Dutch Birding 27: 222, plate 272, 2005).
2004 6 November, Camperduin, *Bergen*, Noord-Holland, adult male (N F van der Ham, S van Kasteel, S Tambach et al).

Reports on the Continental Shelf on 23 November and at Camperduin and Petten, Noord-Holland, on 6 December are still in circulation. The bird at Brouwersdam was possibly the same as the first-winter male present at the same site from 18 January to 7 February 2004 (cf van der Vliet et al 2005) but was accepted as a different bird because, presumably, more than one Surf Scoter can be found in winter in the North Sea area.

Bufflehead / Buffelkopeend
Bucephala albeola 0,0,0
A first-winter male (or more than one bird) at various sites between 5 November 2004 and 18 May 2005, and returning the next winter, remains in circulation to determine the status.

479 Ring-necked Duck / Ringsnaveleend *Aythya collaris*, first-year female, Budel-Dorplein, Noord-Brabant, 23 October 2005 *(René Weenink)* **480** Eastern Imperial Eagle / Keizerarend *Aquila heliaca*, third calendar-year, Kamperhoek, Flevoland, 3 April 2005 *(Ruud F J van Beusekom)* **481** Falcated Duck / Bronskopeend *Anas falcata*, adult male, with Eurasian Wigeon / Smient *A penelope*, male, Nijkerkernauw, Gelderland, 30 March 2005 *(Max Berlijn)* **482** Marbled Duck / Marmereend *Marmaronetta angustirostris*, juvenile, Doornenburg, Gelderland, 15 August 2004 *(Phil Koken)*

Falcated Duck / Bronskopeend
Anas falcata 0,7,2
26 February to 2 March, Roelofsput, *Twenteland*, Overijssel, adult male, photographed (R Gerritsen et al; Dutch Birding 27: 148, plate 184, 2005); 26 March to 5 May, Nuldernauw, *Putten*, Gelderland, adult male, photographed (R O Winters, M Berlijn et al; Birding World 18: 148, 2005, Dutch Birding 27: 222, plate 273, 2005).

The male at Nuldernauw was reportedly seen on the other side of the Nuldernauw at Zeewolde, Flevoland, into June but the CDNA did not (yet) receive details.

American Wigeon / Amerikaanse Smient
Anas americana 5,43,1
3-10 April and 27-30 November, Gaarkeuken, *Zuidhorn*, Groningen, adult male, photographed (S Norrby, R Cazemier, M Bot et al; Dutch Birding 27: 222, plate 274, 2005).
2004 19 November to 1 February 2005, Heerpolderweg, Schelphoek and Middelplaten, *Goes*, Zeeland, first-winter male, photographed.
2002 6-16 April, Kollumerwaard, *Kollumerland en Nieuwkruisland*, Friesland, adult male, photographed (R Cazemier, D Hoekstra, S Bernardus et al).

Note the date extension for the long-staying 2004-05 bird (cf van der Vliet et al 2005). Other reports, eg, near Emmapolder, Groningen, on 18 September and near Sint-Maartensvlotbrug, Noord-Holland, on 2-3 October, have not (yet) been submitted (cf van Dongen et al 2005).

Blue-winged Teal / Blauwvleugeltaling
Anas discors 6,27,0
18-22 April, De Geul and Groote Vlak, *Texel*, Noord-

Holland, adult male, photographed (P Duin, V Stork, J Witte et al).
2004 31 March, Camperduin, *Bergen*, Noord-Holland, adult male (N F van der Ham).
2003 15 April, Camperduin, *Bergen*, Noord-Holland, adult male (N F van der Ham).

The 2005 bird is considered to be the returning 2004 bird that was seen at the same site on Texel (cf van der Vliet et al 2005). Both birds at Camperduin in 2003 and 2004 were fly-bys seen during seawatching and therefore not considered to refer to the same individual. A report on 3 May at Camperduin, Noord-Holland, is still in circulation.

Baikal Teal / Siberische Taling
Anas formosa 8,0,0

The bird collected at Damwoude, Friesland, on 29 November 1948 concerns a female, probably an adult, instead of a first-year male as stated in van IJzendoorn et al (1996) and van den Berg & Bosman (1999, 2001). Re-examination of additional photographs of a presumed Baikal Teal at Broekhuizenvorst, Limburg, from 25 December 1994 to at least 8 April 1995 resulted in its re-identification as a hybrid resembling Baikal Teal (cf Br Birds 88: 269, plate 268, 1995, Dutch Birding 17: 34, plate 31, 1995). This bird had previously been accepted regarding identification but was presumed to have been an escape based on its tame behaviour (Wiegant et al 1996, van den Berg & Bosman 1999, 2001).

*** Marbled Duck / Marmereend**
Marmaronetta angustirostris 0,1,0
2004 14-16 August, Klompenwaard, Doornenburg, *Bemmel*, and Pannerden, *Rijnwaarden*, Gelderland, juvenile, photographed, videoed (R O Winters, H van Diek, P Palmen et al; Winters 2004; Birding World 17: 328, 2004, Dutch Birding 26: 346, plate 491, 357, plate 512-513, 2004).

The first record of a presumed wild bird. Its acceptance was aided by the apparent northward dispersal of (especially) juveniles of this species to north-eastern Spain and southern France in late summer.

Balearic Shearwater / Vale Pijlstormvogel
Puffinus mauretanicus 5,67,-
1996 24 July, Westkapelle, *Veere*, Zeeland (M S J Hoekstein).

This record was previously lost in the mail, so the CDNA is happy that it has finally been accepted. This species is no longer considered since 1 January 1998.

Atlantic Great Cormorant / Grote Aalscholver
Phalacrocorax carbo carbo 6,11,0
2003 27 December, *Heel*, Limburg, male, adult, summer plumage, colour-ring read, photographed (P Palmen).
2002 17 November, *Heel*, Limburg, male, adult, summer plumage, colour-ring read (J Vrehen).
2001 4 November, *Heel*, Limburg, male, adult, summer plumage, colour-ring read (J Vrehen).
2000 4-15 February, *Heel*, Limburg, male, adult, summer plumage, colour-ring read, photographed, sound-recorded (A McGeehan, A B van den Berg, A J van Loon et al; van den Berg et al 2000; Dutch Birding 22: 22, plate 21-22, 2000), and 16 September to 7 January 2001, *Heel*, Limburg (J Smeets, J Vrehen); 26 February, Zuidpier, IJmuiden, *Velsen*, Noord-Holland, immature, photographed (D J Moerbeek); 7 April, Zuidpier, IJmuiden, *Velsen*, Noord-Holland, three, immature, photographed (D J Moerbeek).
1998 6 September, *Heel*, Limburg, second-year, colour-ring read, photographed, sound-recorded (T Cuypers; cf van den Berg et al 2000).

The annually returning Heel individual was colour-ringed (orange with black markings AX on its left leg) as a chick in a colony at Abberton Reservoir near Colchester, Essex, England, on 3 May 1997 (cf van den Berg et al 2000). This is a mixed colony where both (nominate) Atlantic Great Cormorant and Continental Great Cormorants *P c sinensis* breed. Because the identification criteria to separate the two taxa were not yet firmly established at the time, the CDNA decided not to accept it as Atlantic Great (cf van der Vliet et al 2001). Since then, however, Newson et al (2004) published a set of identification criteria, after which the CDNA decided to recirculate and accept the Heel bird, as well as the four IJmuiden birds; the latter four are the first birds without colour-rings to be accepted.

Squacco Heron / Ralreiger
Ardeola ralloides 25 (since 1866),34,2
15 June to 2 July, Oostvaardersplassen, *Almere*, Flevoland, adult, summer plumage, photographed (P Julsing, A Vink et al); 1-2 July, *Alblasserdam*, Zuid-Holland, adult, summer plumage, photographed (L Tukker, C van Rijswijk et al; Dutch Birding 27: 354, plate 432, 2005).
2003 7 June, Quackjeswater, *Westvoorne*, Zuid-Holland, adult, summer plumage (H Janse, W L Janse, B Brieffies et al).

With the additional record for 2003, this is now the best year since 1866 with seven individuals (cf van der Vliet et al 2005).

Glossy Ibis / Zwarte Ibis
Plegadis falcinellus 116,63,-
1935 September, *Zevenaar*, Gelderland, collected (A E M van Nispen; cf Lensink 1993).

The year was erroneously given as 1936 in van den Bergh et al (1979) (Jan van der Brugge in litt, Henk Tamerius in litt). This species is no longer considered since 1 January 2000.

Eurasian Griffon Vulture / Vale Gier
Gyps fulvus 9,54,1
22-23 June, Zuid-Eierland, Oosterend and Polder Waal en Burg, *Texel*, and 23 June, *Den Helder*, Noord-Holland, immature (A Wassink, P Duin, R E Brouwer et al).

Reports at Oosterbeek, Gelderland, on 24 June and at Heerhugowaard, Noord-Holland, on 10 July have not (yet) been submitted (cf van Dongen et al 2005).

Short-toed Eagle / Slangenarend
Circaetus gallicus 3,20,4
17-22 June, Deelensche Veld, *Ede*, Gelderland, photographed (A Vink, W Janssen, D Groenendijk et al; Dutch Birding 27: 284, plate 344, 2005); 28 June, Strabrechtse Heide, *Someren*, Noord-Brabant (R G Bouwman); 2 July to 18 August, Fochteloërveen, *Oost-stellingwerf*, Friesland, and *Norg*, Drenthe, immature, photographed (A de Bruin, R Cazemier, P van Wetter et al; Dutch Birding 27: 356, plate 437-438, 2005); 31 August, De Wijert, *Groningen*, Groningen (A van der Spoel).
2002 20 May, Beusichem, *Amerongen*, Gelderland, immature (A Linckens, W L Janse, F van der Lans et al).

Several reports are still in circulation. Others have not (yet) been submitted (cf van Dongen et al 2005): at Halfweg, Noord-Holland, on 21 May; De Logt, Noord-Brabant, on 22 May; Reeuwijkse Plassen, Zuid-Holland, on 27 May; Marnewaard, Groningen, on 29 May; Naarden, Noord-Holland, on 9 June; Alkmaar, on 26 June; Maastricht, Limburg, on 9 July; Enspijk, Gelderland, on 10 July; Hoge Veluwe, Gelderland, on 12 July; Knardijk, Flevoland, on 23 July; Almere, Flevoland, on 1 August; Lauwersmeer, Groningen, on 17 August; and Dronten, Flevoland, on 13 September.

Pallid Harrier / Steppekiekendief
Circus macrourus 5,19,3
29-30 March, Sint Laurens, *Middelburg*, Zeeland, third calendar-year or adult male, photographed, videoed (A van Gilst, N de Schipper, M Berlijn et al; Dutch Birding 27: 223, plate 276, 2005); 21 April, Breskens, *Oostburg*, Zeeland, second calendar-year male, photographed (M Veldt, I Meulmeester et al; Dutch Birding 27: 223, plate 275, 2005); 24 June, *Alkmaar*, Noord-Holland, adult male (N F van der Ham).

There were no previous records for March and June. Before 2005, records date from 9 April to 17 May in spring and 23 August to 19 October in autumn. Four reports are still in circulation: at Ruurlo, Gelderland on 30 April; Eemshaven, Groningen, on 27 May; Goudzwaard, Zuid-Holland, on 28 August; and Driebergen, Utrecht, on 13 September. Several reports have not (yet) been submitted (cf van Dongen et al 2005): at Posterholt, Limburg, on 26 March; Kamperhoek, Flevoland, on 30 March; Groningen, Groningen, on 4 April; Eemshaven, Groningen, on 11 and another on 25 April; Hoogkerk, Groningen, on 22 April; Sellingen, Groningen, on 2 September; Biggekerke, Zeeland, on 3 September; and Oosternijkerk, Friesland, on 7 September.

Long-legged Buzzard / Arendbuizerd
Buteo rufinus 1,2,0
A bird photographed near Burgh-Haamstede, Zeeland, on 3 October (cf Dutch Birding 27: 431, plate 535, 2005) is still in circulation.

Lesser Spotted Eagle / Schreeuwarend
Aquila pomarina 3,3,1
24-25 September, Domburg, Westkapelle, Oostkapelle and Grijpskerke, *Veere*, Zeeland, juvenile, photographed, videoed (T Luiten, P Beeke, N de Schipper et al; Luiten 2005; Birding World 18: 373, 2005, 19: 25, 2006, Dutch Birding 27: 427, plate 528-530, 2005).

483 Lesser Spotted Eagle / Schreeuwarend *Aquila pomarina*, juvenile, Domburg, Zeeland, 24 September 2005 *(Pim A Wolf)*

Only the second twitchable bird (the other was in November 1984). Found on a Saturday evening, it stayed overnight and could be observed the next day by c 200 birders at close range, mainly in flight. Around noon, it disappeared and was not seen again.

Greater Spotted Eagle / Bastaardarend
Aqula clanga 15,12,0
1957 18 August, Lettele, *Diepenveen*, Overijssel, immature, collected, skin retained at Brabants Natuurmuseum, Tilburg, Noord-Brabant.

This record was erroneously published for 1959 in van der Vliet et al (2005). The museum label indicates that it originates from 1957.

Booted Eagle / Dwergarend
Aquila pennata 0,13,1
20 May, De Hamert, *Arcen en Velden*, Limburg, pale morph, photographed (P Palmen).
2004 1 May, Boshoverheide, *Weert*, Limburg, and Loozerheide, *Cranendonk*, Noord-Brabant, pale morph, videoed (R Verbraak, F Neijts, N van Houtum et al).

The following five reports have not (yet) been submitted (cf van Dongen et al 2005): at Kennemerduinen, Noord-Holland, on 14 May; Hoogkerk, Groningen, on

27 May; Deelensche Veld, Gelderland, on 1 June and another on 15 June; and Veenendaal, Utrecht, on 12 October.

*** Eastern Imperial Eagle / Keizerarend**
Aquila heliaca 0,0,1
3 April, Kamperhoek, *Dronten*, Flevoland, third calendar-year, photographed (R F J van Beusekom, R Alma, M Roos et al; van Beusekom 2005; Birding World 18: 149, 2005, Dutch Birding 27: 233, plate 290-291, 2005).

The bird was observed by seven birders when flying north over the (now) famous migration observation post at Kamperhoek.

Gyr Falcon / Giervalk
Falco rusticolus 4,3,0
2004 13 November, Holwerd, and 19-20 December, Tibma, *Dongeradeel*, Friesland, juvenile, photographed.

Already published in van der Vliet et al (2005) but mentioned here because the CDNA has decided that only one individual was involved.

Baillon's Crake / Kleinst Waterhoen
Porzana pusilla 88,35,6
20-28 May, De Tjamme, Beerta, *Winschoten*, Groningen, singing, photographed, sound-recorded, videoed (B-J Prak, M Bot, W Hartholt et al); 30 May to at least 18 August, Polder Achteraf, Loosdrecht, *Wijdemeren*, Noord-Holland, at least three, adult, singing, photographed, sound-recorded (G van Duin, M Veldt, H R Russer et al; Constantine & The Sound Approach 2006); 3 June to at least 12 August, Nieuwe Keverdijkse Polder, Naardermeer, *Naarden*, Noord-Holland, at least two, singing, sound-recorded (R F J van Beusekom, M S Robb).

Successful breeding was reported at both Polder Achteraf (cf Constantine & The Sound Approach 2006) and Nieuwe Keverdijkse Polder but the outcome has not been submitted to the CDNA, so no young have (yet) been added to the totals. A report at Weerribben, Overijssel, on 18 June, has not (yet) been submitted (cf van Dongen et al 2005). New information indicates that both males and females have an advertisement call, which differs between the sexes. As the ultimate evidence has not been published, the CDNA now considers it unwise to sex recorded birds.

Stone-curlew / Griel
Burhinus oedicnemus -,44,3
24 April, Meijendel, *Wassenaar*, Zuid-Holland (A Remeeus); 2 May, Balgzand, *Anna Paulowna*, Noord-Holland, photographed (R W M Klasens, B Schricken); 6 August, Swifterbant, *Dronten*, Flevoland, first-winter, photographed, videoed (R Alma, O de Vries, M Plomp et al; Dutch Birding 27: 354, plate 431, 2005).

A report on Terschelling, Friesland, on 5 May, has not (yet) been submitted (cf van Dongen et al 2005).

*** Oriental Pratincole / Oosterse Vorkstaartplevier**
Glareola maldivarum 0,1,0
1997 1-5 August, Workumerwaard and Doniaburen, *Wûnseradiel*, Friesland, first-summer, photographed, videoed (M Zekhuis, E Nieuwstraten, G Driessens et al; Nieuwstraten & Zekhuis 1997; Birding World 10: 290, 1997, Dutch Birding 19: 201, plate 204, 211, plate 218-219, 1997).

Although originally identified and submitted as Oriental Pratincole (cf van den Berg & Bosman 1999), it was accepted, after long debates, as an adult Collared Pratincole *G pratincola* (Wiegant et al 1999, van den Berg & Bosman 2001). A record of an Oriental at Ängsnäset, Falsterbo, Skåne, Sweden, from 2 July to 6 October 2001 created renewed interest in the 'Doniaburen pratincole'. Several identification features were critically investigated (cf Driessens & Svensson 2005), resulting in the recirculation of the record. After again long debates and, amongst others, delays by files getting lost in the mail, it was not until 2006 when the bird finally was accepted as Oriental, the first for the Netherlands and, by then, the eighth for the Western Palearctic. A paper discussing this bird will be published in Dutch Birding before long.

*** Killdeer / Killdeerplevier**
Charadrius vociferus 0,0,1
3-5 April, Rottige Meente, *West-Stellingwerf*, Friesland, and Ossenzijl, *IJsselham*, Overijssel, second calendar-year, photographed, sound-recorded, videoed (R-J van der Leij, H Brandsma, E B Ebels et al; van der Leij & Ebels 2005, 2006; Birding World 18: 149, 2005, Dutch Birding 27: 221, plate 270, 223, plate 278, 234, plate 292, 2005, 28: 8, plate 10-11, 2006).

This first for the Netherlands was found on the same day as another first, the Eastern Imperial Eagle at Kamperhoek, Flevoland. The Killdeer was shown on prime time on Dutch television.

Greater Sand Plover / Woestijnplevier
Charadrius leschenaultii 1,7,1
30 April, *Harlingen*, Friesland, summer plumage, photographed (S Bernardus, O Tol, R Cazemier et al; Dutch Birding 27: 223, plate 277, 2005).

Most probably, this was the same bird as the one at Halmstad, Halland, Sweden, on 3-4 May, based on the presence of a single white feather in the left wing. It belonged to the subspecies group *C l leschenaultii/crassirostris*. A record near Battenoord, Zuid-Holland, on 16 September 2004 is still in circulation to determine the subspecies; it has already been accepted and published as Greater Sand Plover (cf van der Vliet et al 2005).

Pacific Golden Plover / Aziatische Goudplevier
Pluvialis fulva 9,22,0

After review, the CDNA has decided that an adult in summer plumage at Middelburg, Zeeland, on 13-15 September 1991 should remain accepted as Pacific Golden Plover.

484 Squacco Heron / Ralreiger *Ardeola ralloides*, adult, Alblasserdam, Zuid-Holland, 2 July 2005 *(Chris van Rijswijk)* **485** Great Snipe / Poelsnip *Gallinago media*, Hessenpoort, Zwolle, Overijssel, 13 September 2005 *(Lennaert Steen)* **486** Eurasian Pygmy Owl / Dwerguil *Glaucidium passerinum*, Sumarreheide, Friesland, 4 October 2002 *(Jan Deinum)* **487** European Roller / Scharrelaar *Coracias garrulus*, adult, Texel, Noord-Holland, 16 September 2005 *(Leo J R Boon/Cursorius)*

Semipalmated Sandpiper / Grijze Strandloper
Calidris pusilla 0,6,1
21-26 August, Wagejot, *Texel*, Noord-Holland, adult, summer plumage, photographed, videoed (A Wassink, D Kok, B van den Boogaard et al; Ebels 2005a; Birding World 18: 321, 2005, Dutch Birding 27: 354, plate 429, 365, plate 445, 2005).

The first record since 1998. The bird was observed well and photographs even show the notorious webbing between the toes! Previous records were in June 1989, July 1996, July 1997 (also at Wagejot, Texel), July 1997, August 1997 and August 1998.

Red-necked Stint / Roodkeelstrandloper
Calidris ruficollis 0,4,1
24 July, Wagejot, *Texel*, Noord-Holland, adult, summer plumage, photographed, videoed (A Wassink, P Duin, R E Brouwer et al; Ebels 2005a; Birding World 18: 277, 2005, Dutch Birding 27: 345, plate 416, 365, plate 446, 2005).

After the untwitchable bird in 2004, birders soon got another chance to see this species. There are now five records with previous ones in May 1987, July 1996, July 1998 and July 2004, all summer-plumaged birds.

White-rumped Sandpiper / Bonapartes Strandloper
Calidris fuscicollis 1,23,3
27 July to 1 August, Scherpenissepolder, *Tholen*, Zeeland, adult, summer plumage moulting to winter plumage, photographed (M Berlijn, G Troost, N de Schipper et al; Dutch Birding 27: 354, plate 430, 2005); 1-2 August, Paesens, *Dongeradeel*, Friesland, adult, summer plumage moulting to winter plumage, photographed (T Bakker, O Tol, R Cazemier et al); 8-9 October, Dijkmanshuizen, *Texel*, Noord-Holland, first-winter, photographed (N van Duivendijk, J de Bruyn, D Kok et al).

Previously, there were only three records later than

488 Buff-breasted Sandpiper / Blonde Ruiter *Tryngites subruficollis*, juvenile, Harpel, Groningen, 2 October 1982 *(Nico de Vries)* **489** Red-necked Stint / Roodkeelstrandloper *Calidris ruficollis*, adult, Wagejot, Texel, Noord-Holland, 24 July 2005 *(René Pop)* **490** Terek Sandpiper / Terekruiter *Xenus cinereus*, Wagejot, Texel, Noord-Holland, 15 May 2005 *(René Pop)* **491** Terek Sandpiper / Terekruiter *Xenus cinereus*, De Braakman, Zeeland, 7 June 2005 *(Mark S J Hoekstein)* **492** Killdeer / Killdeerplevier *Charadrius vociferus*, second calendar-year, Rottige Meente, Overijssel, 4 April 2005 *(Roland Jansen)* **493** Killdeer / Killdeerplevier *Charadrius vociferus*, second calendar-year, Rottige Meente, Overijssel, 4 April 2005 *(Leo J R Boon/Cursorius)*

September: on 18-24 October 1977, 12 October 2000 and 4-5 November 2001.

Broad-billed Sandpiper / Breedbekstrandloper
Limicola falcinellus 23,90,0
2002 16-21 August, Scherpenissepolder, *Tholen*, Zeeland, juvenile, photographed (P A Wolf, M Berlijn et al).
This species is no longer considered since 1 January 2003.

Buff-breasted Sandpiper / Blonde Ruiter
Tryngites subruficollis 3,25,0
2001 15 August, Makkumer Zuidwaard, *Wûnseradiel*, Friesland (M Versluys).
1982 2 October, Harpel, *Vlagtwedde*, Groningen, juvenile, photographed (N de Vries).
The 1982 record is chronologically only the fourth. A report at Zwarte Haan, Friesland, on 5 August has not (yet) been submitted (cf van Dongen et al 2005).

Great Snipe / Poelsnip
Gallinago media -,23,1
13 September, Hessenpoort, *Zwolle*, Overijssel, photographed (D C Uit de Weerd, L J Steen).
September is the best month, with eight records since 1980. Two reports have not (yet) been submitted (cf van Dongen et al 2005): at Kamperhoek, Flevoland, on 28 April, and Nolledijk, Zeeland, on 14 October.

Long-billed Dowitcher / Grote Grijze Snip
Limnodromus scolopaceus 0,19,0
2004 30 September to 18 April 2005, Veerse Meer, Oud-Sabbinge, *Goes*, Zeeland, adult, photographed.
The returning individual already mentioned in van der Vliet et al (2005).

Lesser Yellowlegs / Kleine Geelpootruiter
Tringa flavipes 1,14,1
8-23 October, Prunjepolder, *Schouwen-Duiveland*, Zeeland, adult, winter plumage, photographed, videoed (M Bot, M de Lange et al).
A report from 1964 will be reviewed (again) and a report from 2001 has only recently been submitted.

Terek Sandpiper / Terekruiter
Xenus cinereus 1,30,2
14 May, Dijkmanshuizen, and 15-18 May, Wagejot, *Texel*, Noord-Holland, summer plumage, photographed (A Wassink, A Luijendijk, P Duin et al; Dutch Birding 27: 283, plate 342, 2005); 6-11 June, Braakman Noord, *Terneuzen* (A Kind, P A Wolf, M S J Hoekstein et al), and 4 July, Sophiapolder, *Oostburg*, Zeeland, photographed (P A Wolf, A Klepp, J Eerbeek et al).
Despite the time gap of three weeks, the latter two records are considered to refer to the same bird. A report near Swalmen, Limburg, on 5 May has not (yet) been submitted (cf van Dongen et al 2005).

Franklin's Gull / Franklins Meeuw
Larus pipixcan 0,6,1
13 July, Kwade Hoek, Stellendam, *Goedereede*, Zuid-Holland, adult, summer plumage, photographed (P Stins, P A Wolf et al; Dutch Birding 27: 355, plate 434, 2005).
2004 8 December, Zuidpier, IJmuiden, *Velsen*, Noord-Holland, adult, winter plumage (B van Dijk, B van Buren).
The winter record is not without precedent, as one was seen in February-March 2000. However, midsummer records are the norm.

Kumlien's Gull / Kumliens Meeuw
Larus glaucoides kumlieni 0,0,0
The second-winter photographed on Terschelling, Friesland, on 30 January is still in circulation (Bunskoek 2005; Dutch Birding 27: 153, plate 193-195, 2005).

Ring-billed Gull / Ringsnavelmeeuw
Larus delawarensis 0,8,0
17 September to 13 March 2006, *Tiel*, Gelderland, adult male, photographed (R Gerritsen et al; Dutch Birding 28: 58, plate 77, 2006).
2004 28 February, and 6-13 and 29-30 March, Hoornse Plas, and 16-21 April and 22-25 May, Hoogkerk, *Groningen*, Groningen, first-winter moulting to first-summer, photographed (T Bakker, W Steenge, M Jonker et al; Dutch Birding 26: 214, plate 295, 2004); 4 August to 9 March 2005, *Tiel*, Gelderland, adult male; 27 July to 5 March 2005, *Goes*, Zeeland, adult male.
The Groningen bird is only the second that was not in adult plumage, after a second-winter in April 1996. The Goes bird, first seen in January 1998, did not return in the winter 2005-06.

Ross's Gull / Ross' Meeuw
Rhodostethia rosea 1,13,0
2004 29 November, Camperduin, *Bergen*, Noord-Holland, adult, winter plumage (N F van der Ham).
This is considered to refer to the same bird as present at Scheveningen, Zuid-Holland, on 20-24 and 27 November (van der Vliet et al 2005).

Black Guillemot / Zwarte Zeekoet
Cepphus grylle 31,53,3
16 July, Nes, *Ameland*, Friesland, adult, summer plumage (M Bot); 20 August, Maasvlakte, *Rotterdam*, Zuid-Holland, adult, summer plumage (M Renders, N van Houtum); 6-15 October, Westerslag, and 18-19 October, De Koog, *Texel*, Noord-Holland, adult, winter plumage, photographed (P L Meininger et al).
2004 29 July, Camperduin, *Bergen*, Noord-Holland, first-winter (N F van der Ham); 20 September to 24 March 2005, West-Terschelling, *Terschelling*, Friesland, adult, winter plumage moulting to summer plumage, photographed.
2003 6 February, Camperduin, *Bergen*, Noord-Holland, adult, winter plumage (N F van der Ham, G J de Haan).
2002 20 October, Camperduin, *Bergen*, Noord-Holland, adult, winter plumage (M Platteeuw, T Knol, N F van der Ham); 11 November, Camperduin, *Bergen*, Noord-Holland, adult, winter plumage (N F van der Ham, E Dekker).



494 Northern Hawk Owl / Sperweruil *Surnia ulula*, first-year, Westerbork, Drenthe, 31 October 2005 *(Roland Jansen)* **495** Northern Hawk Owl / Sperweruil *Surnia ulula*, first-year, Westerbork, Drenthe, 31 October 2005 *(Harvey van Diek)* **496** Thrush Nightingale / Noordse Nachtegaal *Luscinia luscinia*, Flevocentrale, Flevoland, 21 May 2005 *(Roland Jansen)* **497** Greater Short-toed Lark / Kortteenleeuwerik *Calandrella brachydactyla*, Kennemerduinen, Bloemendaal, Noord-Holland, 18 November 2005 *(Arnoud B van den Berg/Vrs van Lennep)*

2000 30 November, Camperduin, *Bergen*, Noord-Holland, adult, winter plumage (N F van der Ham, G J de Haan).

This species will no longer be considered from 1 January 2007 onwards.

Eurasian Eagle Owl / Oehoe
Bubo bubo 1,45,-
1992 1 October, Mook, *Mook en Middelaar*, Limburg, photographed, picked up injured and taken into care at Burgers' Dierenpark, Arnhem, Gelderland (M van Ginneke).

This record was lost in the mail, so the CDNA is pleased that it has finally been accepted. It was chronologically only the fifth record. This species is no longer considered since 1 January 2004.

Northern Hawk Owl / Sperweruil
Surnia ulula 1,1,1
30-31 October, Hooghalen, *Aa en Hunze*, Drenthe, first-winter, photographed, videoed (H L Holtkamp, E J Alblas, P van Wetter et al; Alblas 2005, van Wetter et al 2006; Birding World 18: 425, 2005, 19: 31, 2006, Dutch Birding 27: 429, plate 533, 450, plate 562, 2005, 28: 292, plate 409, 293, plate 410, 294, plate 411-413, 2006).

For most birders *the* highlight of 2005. It showed very well on its second day (when the news broke) but surprisingly had left the area the following day. It belonged to the nominate subspecies *S u ulula*. Previous records were on 5 October 1920 at Amerongen, Utrecht (collected) and 2 April 1995 at Brunssum, Limburg.

*** Eurasian Pygmy Owl / Dwerguil**
Glaucidium passerinum 0,1,0
2002 4 October, Sumarreheide, Sumar (Suameer), *Tytsjerksteradiel*, Friesland, picked up and released, videoed (J Deinum; de Jong 2004ab).

The first record. Because there were no reports of notable southward movements in Scandinavia in the autumn of 2002, this bird most probably originated from the German population; the nearest breeding locations are in Nordrhein-Westfalen and Niedersachsen. It was first identified as Little Owl *Athene noctua* and only correctly reidentified in February 2004, when video-stills were sent to Johan de Jong.

Alpine Swift / Alpengierzwaluw
Apus melba 7,37,2
30 April, Breskens, *Oostburg*, Zeeland, photographed (G Troost, J Leclercq; Dutch Birding 27: 284, plate 343, 2005); 6 May, Huizer Pier, *Huizen*, and Stichtse Brug, *Blaricum*, Noord-Holland (H van Oosterhout, R Alma).

European Roller / Scharrelaar
Coracias garrulus 53,16,1
15-20 September, Ecomare, *Texel*, Noord-Holland, adult, photographed (L J R Boon, R van Rossum et al; Birding World 18: 372, 2005, Dutch Birding 27: 350, plate 424, 429, plate 534, 431, plate 536, 2005).

The first since 1998, reflecting an obvious decline in records. The best months are May and June with 21 and 27 records, respectively. There are now five records from September, and, coincidentally, also five from Texel, with previous records in May 1964, June 1966, May 1971 and June 1980. A report near Huijbergen, Noord-Brabant, on 29 May, has not (yet) been submitted (cf van Dongen et al 2005).

Calandra Lark / Kalanderleeuwerik
Melanocorypha calandra 0,2,1
27 May, Eemshaven-Oost, Eemshaven, *Eemsmond*, Groningen, photographed (R Hovinga, E Klunder, M Bot et al; Hovinga & Olthoff 2005, Hovinga 2006; Dutch Birding 27: 294, plate 360, 2005, 28: 231, plate 314, 2006).

This bird stayed for c 1.5 h. Previous records were at Castricum, Noord-Holland, on 10 October 1980 (ringed), and on Texel, Noord-Holland, on 16 May 1988.

Greater Short-toed Lark / Kortteenleeuwerik
Calandrella brachydactyla 3,29,2
21 April, Hee, *Terschelling*, Friesland, photographed (A Ouwerkerk; Birding World 18: 148, 2005, Dutch Birding 27: 225, plate 282, 2005); 18 November, Kennemerduinen, *Bloemendaal*, Noord-Holland, ringed, photographed (J Vegelin, J van Tussenbroek, P Kooij et al).

As in previous years, there were several records of fly-bys at different migration hot-spots which have not been submitted. One briefly reported at Westerplas, Schiermonnikoog, Friesland, on 25 September has not (yet) been submitted either. The Kennemerduinen bird is the latest ever; there are only two previous November records (7 November 1981 and 10 November 2000).

Olive-backed Pipit / Siberische Boompieper
Anthus hodgsoni 0,15,1
7 October, Maasvlakte, *Rotterdam*, Zuid-Holland, photographed, videoed (W van Rijswijk, J den Hertog, E B Ebels et al; Dutch Birding 27: 431, plate 538, 2005).
2004 3-6 October, Lange Paal, *Vlieland*, Friesland (F Ossendorp, H A M Prinsen, L B Steijn et al).

The 2004 bird is another individual than the one already accepted for exactly the same four days near the Badweg on the same island (cf van der Vliet et al 2005). A report on Texel, Noord-Holland, on 13 October is still in circulation (cf van Dongen et al 2005).

Red-throated Pipit / Roodkeelpieper
Anthus cervinus 25,51,-
1981 10-11 May, Rijswijkse Buitenpolder, *Buren*, Gelderland, maximum of two (on 11 May).

This record has previously been published with wrong municipality and province (cf van den Berg & Bosman 1999, 2001). This species is no longer considered since 1 January 1992.

Thrush Nightingale / Noordse Nachtegaal
Luscinia luscinia 7,49,3
21 May into June, Flevocentrale, *Lelystad*, Flevoland, at least two (second from at least 25 May), at nest, sing-

498 Desert Wheatear / Woestijntapuit *Oenanthe deserti*, first-year male, IJmuiden, Noord-Holland, 24 September 2005 *(René van Rossum)*

499 Desert Wheatear / Woestijntapuit *Oenanthe deserti*, first-year male, IJmuiden, Noord-Holland, 22 November 2005 *(Leo J R Boon/Cursorius)*

ing, photographed, sound-recorded (G J ter Haar, R Halff, F Geldermans et al; Dutch Birding 27: 286, plate 348, 2005); 22-25 May, Meyendel, *Wassenaar*, Zuid-Holland, singing, sound-recorded (J Noordhuizen, R van der Vliet, A Remeeus et al).

Breeding was reported at Flevocentrale but the outcome has not been submitted to the CDNA. This would represent the second breeding record; the first breeding took place in 1995 at Horsterwold, Flevoland (van Beusekom 2001).

Red-spotted Bluethroat / Roodsterblauwborst
Luscinia svecica svecica 24,40,0
2004 9 May, De Cocksdorp, *Texel*, Noord-Holland, male (A Wassink, P Duin); 28 May, De Cocksdorp, *Texel*, Noord-Holland, male (A Wassink, K Ophof).

White-throated Robin / Perzische Roodborst
Irania gutturalis 0,3,1
31 October, Roptazijl, Pietersbierum, *Franekeradeel*, Friesland, first-winter, ringed, found dead on 2 November, skin not retained, photographed (U Rijpma, A Bakker, S van Lingen et al; Rijpma & Bakker 2006; Dutch Birding 27: 434, plate 545, 2005, 28: 302, plate 418, 303, plate 419-420, 2006).

The fourth record and the first of the less well-known pale morph; previous records were at Maasland, Zuid-Holland, on 3-4 November 1986 (male), Wassenaar, Zuid-Holland, on 2 June 1995 (female), and Lozerheide, Limburg, on 30 August 2003 (first-year).

Ehrenberg's Redstart / Oosterse Gekraagde Roodstaart
Phoenicurus phoenicurus samamisicus 0,0,0

A male photographed at Groote Vlak, Texel, Noord-Holland, on 10 September is still in circulation; another male on the same date at Lentevreugd, Texel, has not yet been submitted (cf Wassink & Ebels 2005; Dutch Birding 27: 366, plate 447, 2005).

Siberian Stonechat / Aziatische Roodborsttapuit
Saxicola maurus 9,22,1
26 October, Westduinpark, *Den Haag*, Zuid-Holland, first-winter, photographed (V van der Spek, R Berkelder, R ter Ellen et al; Dutch Birding 27: 436, plate 551, 2005).

Isabelline Wheatear / Izabeltapuit
Oenanthe isabellina 0,3,1
18 November, Buren, *Terschelling*, Friesland, first-winter, photographed (J Vink, A Ouwerkerk; Birding World 18: 454, 2006, Dutch Birding 28: 61, plate 81, 2006).

The fourth record, concerning a bird seen on the North Sea beach for c 1 h, just before dark. Previous records were at Maasvlakte, Zuid-Holland, from 21 October to 8 November 1996, at IJmuiden, Noord-Holland, on 22-23 September 2000 and on Schiermonikkoog, Friesland, on 14-25 October 2000.

Desert Wheatear / Woestijntapuit
Oenanthe deserti 1,9,9
5-10 April, Fochteloërveen, *Norg*, Drenthe, first-summer male, singing, photographed (W Kramer, E Schoppers, R Jansen et al; Birding World 18: 150, 2005, Dutch Birding 27: 226, plate 283, 2005); 23-29 September, Zuidpier, IJmuiden, *Velsen*, Noord-Holland, first-winter male, photographed, videoed (R Rotscheid, T Nagtegaal, R van Rossum et al; Birding World 18: 373, 2005, Dutch Birding 27: 432, plate 541, 2005); 29 October, Paal 2, *Schiermonnikoog*, Friesland, first-winter male, photographed (J Bosma et al; Dutch Birding 27: 435, plate 549, 2005); 30 October to 5 November, Wierum, *Dongeradeel*, Friesland, first-winter female, photographed (H van Kampen, R Cazemier et al; Dutch Birding 27: 432, plate 542, 2005); 5 November, Dashorstdijk, *Almere*, Flevoland, first-winter male, photographed (G Heester, D Roos, P Bertens et al); 7 November, Nes, *Terschelling*, Friesland, first-winter female, photographed (H Goodijk); 11-24 November, Zuidpier, IJmuiden, *Velsen*, Noord-Holland, first-winter male, photographed (R Rotscheid, R Altenburg, R van Dijk et al; Dutch Birding 28: 60, plate 78, 2006); 19 November, Dam 20, *Vlieland*, Friesland, first-winter male, photographed (C Giljam, J Reneerkens, G van Duin et al; Dutch Birding 28: 60, plate 79, 2006); 2 December, De Hon, *Ameland*, Friesland, male, photographed (J Huizenga, IJ de Jong).

In autumn, a remarkable influx of eight individuals took place. The birds at Zuidpier are considered two individuals because this is a well-watched site where a single long-staying bird would not have gone unnoticed for six weeks. The Fochteloërveen bird was the second in spring, after a female in April 1989.

White's Thrush / Goudlijster
Zoothera aurea 11,6,1
29 October, Hasseltsestraat, Scheveningen, *Den Haag*, Zuid-Holland, found dead, photographed (E Kok).

Pallas's Grasshopper Warbler / Siberische Sprinkhaanzanger
Locustella certhiola 0,5,1
20 September, *Castricum*, Noord-Holland, ringed, photographed, videoed (H Levering, A J van Loon, P Veldt et al; Roselaar et al 2006b; Dutch Birding 28: 276, plate 400, 2006).

The fifth record for the ringing site at Castricum; the first was on 5 October 1991. Roselaar et al (2006b) presented an overview of the five September records in 2002-05, also discussing the subspecies.

River Warbler / Krekelzanger
Locustella fluviatilis 6,36,1
1-3 July, Dordtse Biesbosch, *Dordrecht*, Zuid-Holland, singing, photographed (H Gebuis, T O V Muusse, P A Wolf).

Reports at Ooijpolder, Gelderland, on 15 June, and near Taarloo, Drenthe, on 26 July, have not (yet) been submitted (cf van Dongen et al 2005).

Melodious Warbler / Orpheusspotvogel
Hippolais polyglotta 2,45,-
2004 11-16 June, Sint Geertruid, *Margraten*, Limburg, singing, photographed.

This record was published with the wrong municipality in van der Vliet et al (2005). This species is no longer considered since 1 January 2005; de Bruin & van der Vliet (2005) erroneously stated that it was no longer considered since 1 January 2006.

Paddyfield Warbler / Veldrietzanger
Acrocephalus agricola 1,13,1
11 September, Wormer- en Jisperveld, *Waterland*, Noord-Holland, adult, ringed, photographed (J van der Geld, R Leurs; Dutch Birding 27: 434, plate 546, 2005).

Again a bird trapped during ringing activities. There is still only one field record, on Vlieland, Friesland, on 18 September 1994.

Blyth's Reed Warbler / Struikrietzanger
Acrocephalus dumetorum 0,7,0
2003 28-30 September, Stortemelk, *Vlieland*, Friesland, photographed, sound-recorded (J de Bruijn, N van Duivendijk, H Zevenhuizen et al; Dutch Birding 25: 430, plate 503, 2003).

The third for 2003, making it an even better year for this species.

Dartford Warbler / Provençaalse Grasmus
Sylvia undata 1,6,0
2004 29 December to 12 February 2005, Dwingelderveld, *Westerveld*, Drenthe, adult male, photographed, sound-recorded.

This record was already mentioned in van der Vliet et al (2005).

Subalpine Warbler / Baardgrasmus
Sylvia cantillans 5,43,3
10 May, Bakkum, *Castricum*, Noord-Holland, male, photographed (C van der Wardt, M van der Wardt, R Halff et al); 31 October to 1 November, Westerplas, *Schiermonnikoog*, Friesland, female (H Dufourny, V Buchet).
2004 28 May, De Uithof, *Den Haag*, Zuid-Holland, male, photographed, sound-recorded (W Kolber, T J C Luijendijk, F Ossendorp et al).
2003 26-27 April, Westkapelle, *Veere*, Zeeland, first-summer male, photographed (G van de Maat, M Renders, A Linckens et al);
2000 9 May, Rottumerplaat, *Eemsmond*, Groningen, female, photographed (D Lutterop, G Kasemir).

Totals include birds identified to subspecies (see below). The CDNA is still undecided on the subspecific identity of the 2004 bird: for that reason it remains in circulation. A report at Callantsoog, Noord-Holland, on 2 May, has not (yet) been submitted (cf van Dongen et al 2005).

Western Subalpine Warbler / Westelijke Baardgrasmus
Sylvia cantillans cantillans/inornata 2,4,1
7-10 July, Bleekersvallei, *Texel*, Noord-Holland, first-summer male, photographed (K Ophof, A Wassink, J J F J Jansen et al; Dutch Birding 27: 355, plate 436, 2005).
2003 18 May, Kollumeroord, *Kollumerland en Nieuwkruisland*, Friesland, first-summer male, sound-recorded (R Cazemier, M P Olthof, T Bakker et al).

*** Western Orphean Warbler / Westelijke Orpheusgrasmus**
Sylvia hortensis 0,1,0
2003 29 October to 5 November, Arnestein I, *Middelburg*, Zeeland, photographed, videoed (P A Wolf, P L Meininger, M S J Hoekstein et al; Wolf 2003; Birding World 16: 420, 2003, 17: 37, 2004, Birdwatch 12 nr 136: 63, 2003, Dutch Birding 25: 420, plate 490, 444, plate 522, 2003, 26: 71, plate 126, 2004).

A paper discussing the identification (including the separation from Eastern Orphean Warbler *S crassirostris*) has been prepared for Dutch Birding.

Greenish Warbler / Grauwe Fitis
Phylloscopus trochiloides 6,38,3
4-12 June, *Bodegraven*, Zuid-Holland, singing, photographed, sound-recorded (K Jongeneel, T J C Luijendijk, S Schilperoort et al; Dutch Birding 27: 287, plate 351, 2005); 11-12 June, 't Zand, *Zijpe*, Noord-Holland, singing, photographed, sound-recorded (R de Beer, R E Brouwer, A van de Ende et al); 31 August, De Cocksdorp, *Texel*, Noord-Holland (W van Rijswijk, A Wassink et al).
2004 5-9 June, Sint Laurens, *Middelburg*, Zeeland, singing, photographed, sound-recorded (P Beeke, C Beeke, J Walhout et al); 9 August, Groene Glop, *Schiermonnikoog*, Friesland, first-winter, ringed, photographed (A G Duiven, K H Oosterbeek); 3-7 October, Robbenjager, De Cocksdorp, *Texel*, Noord-Holland, photographed, sound-recorded (W L Janse, P van Wetter et al).
2003 29 May, Quackgors, *Hellevoetsluis*, Zuid-Holland, singing, sound-recorded (M Klemann).

A report from Eemshaven, Groningen, on 17-20 and 25 September, is still in circulation (cf van Dongen et al 2005). The record of October 2004 concerns a date extension (cf van der Vliet et al 2005). The record on Schiermonnikoog in early August 2004 may suggest local breeding (especially since the first breeding record was on the same island in 2003, cf Ebels 2003) but there were no other records on the island in 2004 and, therefore, it concerned most likely 'just' an early migrant.

Hume's Leaf Warbler / Humes Bladkoning
Phylloscopus humei 2,28,1
20 November to 14 January 2006, Oosterbeek, *Renkum*, Gelderland, photographed, sound-recorded (S Terlouw, M van Duijn, M P Lantsheer et al).

Radde's Warbler / Raddes Boszanger
Phylloscopus schwarzi 2,14,1
15 October, Oost-Vlieland, *Vlieland*, Friesland (L B Steijn, R F J van Beusekom, N van Duivendijk et al).

Dusky Warbler / Bruine Boszanger
Phylloscopus fuscatus 1,42,1
29 October, *Castricum*, Noord-Holland, ringed, photo-

500 Western Orphean Warbler / Westelijke Orpheusgrasmus *Sylvia hortensis*, Middelburg, Zeeeland, 31 October 2003 *(Bas van den Boogaard)* **501** Desert Wheatear / Woestijntapuit *Oenanthe deserti*, first-summer male, Fochteloërveen, Drenthe, 6 April 2005 *(Roland Jansen)* **502** Two-barred Crossbill / Witbandkruisbek *Loxia leucoptera bifasciata*, male, Wateren, Drenthe, 9 February 2005 *(Roland Jansen)*

graphed (A Wijker et al).
2004 9 October, Ouddorp, *Goedereede*, Zuid-Holland, photographed, sound-recorded (M Renders, N van Houtum, G Bakker et al); 17 October, Oost-Vlieland, *Vlieland*, Friesland (F Ossendorp, J Scharringa, J Eerdmans et al).

The Ouddorp bird was submitted as Radde's Warbler *P schwarzi* but the documentation convinced the CDNA to accept it as Dusky, again showing that the identification of these species is not always straightforward. A report from Oostkapelle, Zeeland, on 22 November has not (yet) been submitted (cf van Dongen et al 2006).

Siberian Chiffchaff / Siberische Tjiftjaf
Phylloscopus collybita tristis -,19,1
11-29 October, Westkapelle, *Veere*, Zeeland, photographed, sound-recorded (P Palmen, P A Wolf et al).
2003 c 15 November to 12 February 2004, *Harlingen*, Friesland, sound-recorded (M Versluys, M S Robb, M Brandsma et al).

Several reports are still in circulation: on Ameland, Friesland, on 19-20 November; at Lauwersmeer, Groningen, from 20 November to 4 December; Kollumerpomp, Friesland, on 26 November to 8 December; and Alkmaar, Noord-Holland, on 23-28 December. The identification of birds at Groene Glop, Schiermonnikoog, Friesland, on 2 November 1995 (trapped) and at Laakse Strand, Zeewolde, Flevoland, on 18 October 1998 have been confirmed after review. One at Schelle, Zwolle, Overijssel, on 20-27 November 1988 has been rejected after review. Two other old records are still being reviewed.

Iberian Chiffchaff / Iberische Tjiftjaf
Phylloscopus ibericus 2,14,1
19 May to 12 June, *Bergen*, Noord-Holland, singing, photographed, sound-recorded, videoed (D Vlugt, B van Wees, R E Brouwer et al; Dutch Birding 27: 284, plate 346, 2005).
2004 24 April to 6 June, Vinkhuizen, *Groningen*, Groningen, male, singing, photographed, sound-recorded (D M Brinkhuizen, A de Bruin, E B Ebels et al).

Collared Flycatcher / Withalsvliegenvanger
Ficedula albicollis 24,8,0
1915 5 April, *Loosdrecht*, Utrecht, adult male (K Boedijn; De Levende Natuur 19 (24): 520, 1915).

Turkestan/Daurian Shrike / Turkestaanse/Daurische Klauwier
Lanius phoenicuroides/isabellinus 0,2,0

After review, the CDNA remains of the opinion that, on present knowledge, it is not possible to decide whether a bird at De Hoge Berg, Texel, Noord-Holland, on 21 October 1993 was a Turkestan Shrike or Daurian Shrike.

Lesser Grey Shrike / Kleine Klapekster
Lanius minor 21,17,1
5 October, Groote Vlak, *Texel*, Noord-Holland, adult female, photographed (K de Mooij, W L Janse, R van Viegen et al; Dutch Birding 27: 436, plate 550, 2005).

Texel is the best place to see this shrike, with six previous records, in June 1964, May 1970, June 1971, May 1978, July 1984 and June 1998.

Red-eyed Vireo / Roodoogvireo
Vireo olivaceus 0,7,1
9 October, Westkapelle, *Veere*, Zeeland, photographed (P L Meininger, C van Rijswijk et al; Meininger 2005; Dutch Birding 27: 435, plate 547-548, 2005).

Previous records were between 24 September and 22 October in 1985 (2), 1991 (2), 1996, 2001 and 2003. This was the second field record, after one in October 1996.

Arctic Redpoll / Witstuitbarmsijs
Carduelis hornemanni 43,57,3
18 November, Zuidpier, IJmuiden, *Velsen*, Noord-Holland, male, photographed (J van 't Hof); 19 November, Oosterscheldekering, Westenschouwen, *Schouwen-Duiveland*, Zeeland, first-winter, photographed (C van Rijswijk); 27 December to 7 January 2006, Marshaven, *Zutphen*, Overijssel, male, photographed, videoed (H van Oosten, J Voerknecht et al; Dutch Birding 28: 63, plate 84-85, 2006).

Several reports are still in circulation. Totals include Hornemann's Redpoll *C h hornemanni* (see below).

*** Hornemann's Redpoll / Groenlandse Witstuitbarmsijs**
Carduelis hornemanni hornemanni 0,1,0
2003 11-15 October, Huisduinen, *Den Helder*, Noord-Holland, male, adult, photographed, sound-recorded, videoed (K Rebel, R Halff, R E Brouwer et al; cf Ebels & Halff 2003; Birding World 16: 455, 2003, Dutch Birding 25: 430, plate 507, 440, plate 519, 2003).

The first record of this taxon. Its occurrence coincided with at least 10 individuals in Scotland between 22 September and 7 October (Rogers et al 2004, 2005). The identification of recent birds in Britain was discussed by Pennington & Maher (2005). A paper detailing the Dutch record will be published in Dutch Birding before long.

Two-barred Crossbill / Witbandkruisbek
Loxia leucoptera bifasciata 36,219,1
30 January to 11 February, Wateren, *Noordenveld*, Drenthe, adult male, photographed (O Tol, R Jansen, M Bot et al; Dutch Birding 27: 147, plate 182-183, 2005).
2002 17 February, Bakkum, *Castricum*, Noord-Holland, adult female (N F van der Ham).

Trumpeter Finch / Woestijnvink
Bucanetes githagineus 0,1,1
12 June, Eemshaven-Oost, Eemshaven, *Eemsmond*, Groningen, male, photographed, videoed (M P Olthoff, T Bakker, M Bot et al; Hovinga & Olthoff 2005, Olthoff 2006; Dutch Birding 27: 294, plate 361, 2005, 28: 235, plate 323-324, 2006).

This second came soon after the first at Maasvlakte,

503 Lesser Grey Shrike / Kleine Klapekster *Lanius minor*, adult female, Groote Vlak, Texel, Noord-Holland, 5 October *(Leon Boon)* **504** Desert Wheatear / Woestijntapuit *Oenanthe deserti*, first-year male, Westerstrand, Schiermonnikoog, Friesland, 29 October 2005 *(Ted Post)* **505-506** Desert Wheatear / Woestijntapuit *Oenanthe deserti*, first-year male, IJmuiden, Noord-Holland, 13 November 2005 *(Karel A Mauer)* **507** Arctic Redpoll / Witstuitbarmsijs *Carduelis hornemanni exilipes*, first-winter, Oosterscheldekering, Westenschouwen, Zeeland, 19 November 2005 *(Chris van Rijswijk)* **508** Hornemann's Redpoll / Groenlandse Witstuitbarmsijs *Carduelis hornemanni hornemanni*, adult male, Huisduinen, Noord-Holland, 15 October 2003 *(Arnoud B van den Berg)*

509 Greenish Warbler / Grauwe Fitis *Phylloscopus trochiloides*, Bodegraven, Zuid-Holland, 10 June 2005 *(Jan den Hertog)* **510** Iberian Chiffchaff / Iberische Tjiftjaf *Phylloscopus ibericus*, Bergen, Noord-Holland, 25 May 2005 *(Leo J R Boon/Cursorius)* **511** Siberian Chiffchaff / Siberische Tjiftjaf *Phylloscopus collybita tristis*, Westkapelle, Zeeland, 11 October 2005 *(Patrick Palmen)* **512** Red-eyed Vireo / Roodoogvireo *Vireo olivaceus*, Westkapelle, Zeeland, 9 October 2005 *(Chris van Rijswijk)*

Zuid-Holland, on 31 May 2003 (van Rijswijk 2004); it was part of an unprecedented influx to north-western and northern Europe (cf Olthoff 2006).

Pine Bunting / Witkopgors
Emberiza leucocephalos 15,18,1
13-16 March, Gastel, *Cranendonk*, Noord-Brabant, female, photographed (R Verbraak, F Neijts, R G Bouwman et al).
2004 1 November, Westenschouwen, *Schouwen-Duiveland*, Zeeland, first-winter female, ringed, photographed (C Oskam, L van Ree).

Rustic Bunting / Bosgors
Emberiza rustica 14,56,1
16 October, Lange Paal, *Vlieland*, Friesland, adult male, photographed (L B Steijn, N van Duivendijk et al; Dutch Birding 27: 436, plate 552, 2005).

Black-headed Bunting / Zwartkopgors
Emberiza melanocephala 2,8,1
20 June, Schinkelpolder, Amsterdamse Bos, *Amstelveen*, Noord-Holland, female, found dead, photographed, skin retained at Zoölogisch Museum Amsterdam (D Kars).

The first female. Previous records were males in May 1962, February 1967, June 1993, May 1996 (two), May 1997, July 1997, June 1998, June-July 2000 and June 2003, making May-June the best period with now nine.

Systematic list of rejected reports
This list contains all reports not accepted by the CDNA. Reports marked with @ were rejected by all committee members in the first or second voting. Most were rejected because the identification was not fully established (often due to lack of documentation) or when the bird showed signs of captivity, such as dubious rings, excessive wear and/or aberrant beha-

viour (for presumed or proven escapes, the text indicates if the identification has been accepted or not). Species for which the CDNA unanimously decided that all records up to now refer to birds of captive origin are indicated with *. This includes records of birds of which all ancestors are assumed to be of captive origin (introduced species of British category C).

Whistling Swan / Fluitzwaan *Cygnus columbianus* 5 February, Vereenigde Binnenpolder, *Haarlemmerliede en Spaarnwoude*, Noord-Holland, adult (@ exact amount of yellow on bill not determined). **Greenland White-fronted Goose / Groenlandse Kolgans** *Anser albifrons flavirostris* 9 January, Someren-Heide, *Someren*, Noord-Brabant, two, immature, photographed (@ description incomplete; photographs do not show conclusive fieldmarks). **Hutchins's Canada Goose / Hutchins' Canadese Gans** *Branta hutchinsii hutchinsii* 26-28 March, *Olst*, Overijssel, photographed (description wrong in some respects, eg, legs too long and breast too dark). * **Bufflehead / Buffelkopeend** *Bucephala albeola* 3 July, Bovenwater, *Lelystad*, Flevoland, female, photographed (@ identification accepted; uncertain whether unringed). **Baikal Teal / Siberische Taling** *Anas formosa* 28 March, De Regulieren, *Geldermalsen*, Gelderland, adult male (@ poor description). **Great White Pelican / Roze Pelikaan** *Pelecanus onocrotalus* 29 April, *Wageningen*, Gelderland, adult (@ description incomplete). **Black-shouldered Kite / Grijze Wouw** *Elanus caeruleus* 13 March, Vaassen, *Epe*, Gelderland, adult (@ description incomplete). **Short-toed Eagle / Slangenarend** *Circaetus gallicus* 21 May, Starrevaart, *Leidschendam*, Zuid-Holland (@ poor description); 10 July, Loozerheide, *Weert*, Limburg (@ description incomplete). **Eurasian Black Vulture / Monniksgier** *Aegypius monachus* 16-17 March, Beers, *Cuijk*, Noord-Brabant, and 20 March to 15 August, Oostvaardersplassen, *Almere/Lelystad*, Flevoland, adult female ('Carmen'), photographed, videoed (found dead on 15 August), skin retained at Nationaal Natuurhistorisch Museum/Naturalis, Leiden, Zuid-Holland (@ identification accepted; the CDNA decided that its occurrence after a translocation of c 1000 km (from central Spain to southern France) can no longer be considered 'natural'; Ebels 2005b; Birding World 18: 100, 2005, Dutch Birding 27: 224, plate 279, 329, plate 403, 330, plate 404, 331, plate 405, 332, plate 406, 2005). **Pallid Harrier / Steppekiekendief** *Circus macrourus* 19-20 April, Rottumeroog, *Eemsmond*, Groningen, female, photographed (@ no description; photographs only show a rather distant head); 1 May, Breskens, *Oostburg*, Zeeland, male, photographed, videoed (@ no description; photographs too blurry); 6 September, Blauwe Kamer, *Rhenen*, Utrecht, juvenile (@ description incomplete). **Golden Eagle / Steenarend** *Aquila chrysaetos* 3 November, Appelscha, *Oststellingwerf*, Friesland, immature (@ poor description). **Booted Eagle / Dwergarend** *Aquila pennata* 20 April, Epser Waarden, Wilp, *Voorst*, Gelderland, pale morph (@ description incomplete); 28 May, Buurserzand, *Haaksbergen*, Overijssel, pale morph (@ description incomplete); 29 May, Bargerveen, *Emmen*, Drenthe, dark morph (@ description incomplete). **Spotted Sandpiper / Amerikaanse Oeverloper** *Actitis macularius* 15 August, Polder Breebaart, *Delfzijl*, Groningen, adult, summer plumage, photographed (@ photographs show Common Sandpiper *A hypoleucos*). **European Roller / Scharrelaar** *Coracias garrulus* 20 September, between Warns and Hemelum, *Nijefurd*, Friesland (@ description indicates that Abyssinian Roller *C abyssinica* was involved, of which an escaped individual is known to have been around in that area at the time). **Grey-headed Woodpecker / Grijskopspecht** *Picus canus* 3 September, Linne, *Maasbracht*, Limburg, adult female (@ description too brief of a bird seen for only 5 sec). **Iberian Wagtail / Iberische Kwikstaart** *Motacilla iberiae* 7 May to 5 June, *Nieuwerkerk aan den IJssel*, Zuid-Holland, male, photographed (@ photographs show first-summer Blue-headed Wagtail *M flava*; Dutch Birding 27: 287, plate 352, 2005). **Black-headed Wagtail / Balkankwikstaart** *Motacilla feldegg* 1 May, *Scherpenzeel*, Gelderland, adult male (@ records of this species can only be accepted if photographs and/or video-recordings are available). **Siberian Thrush / Siberische Lijster** *Zoothera sibirica* 12 October, Sintjohannesga, *Skarsterlân*, Friesland, adult male (@ description partly wrong for this species, eg, size described as larger than Eurasian Blackbird *Turdus merula*). **Siberian Chiffchaff / Siberische Tjiftjaf** *Phylloscopus collybita tristis* 23-24 April, Lauwersoog, *De Marne*, Groningen, sound-recorded (@ description partly wrong for this taxon; sound-recording indicates Common Chiffchaff *P c collybita*). **Arctic Redpoll / Witstuitbarmsijs** *Carduelis hornemanni* 13-14 November, Loenermark, *Apeldoorn*, Gelderland, adult male (@ description incomplete); 22 November, De Bijland, Lobith, *Rijnwaarden*, Gelderland, adult male, ringed, photographed (@ photographs show adult male Mealy Redpoll *C flammea*). **Parrot Crossbill / Grote Kruisbek** *Loxia pytyopsittacus* 6 February, *Wageningen*, Gelderland, adult male (@ description incomplete). **Pine Grosbeak / Haakbek** *Pinicola enucleator* 27 February, *Heemskerk*, Noord-Holland, adult male (@ description incomplete, cf Bosma et al 2005). **Pine Bunting / Witkopgors** *Emberiza leucocephalos* 28 October, Malden, *Heumen*, Gelderland (@ no bird seen; description of only call insufficient for acceptance).

2004

Swinhoe's Storm-petrel / Chinees Stormvogeltje *Oceanodroma monorhis* 22 September, Camperduin, *Bergen*, Noord-Holland (@ description incomplete). **Booted Eagle / Dwergarend** *Aquila pennata* 27 April, Slikken van Flakkee, *Dirksland*, Zuid-Holland, pale morph, photographed (description incomplete; photographs too blurry to show conclusive details). **Pallid Swift / Vale Gierzwaluw** *Apus pallidus* 21 October, Oosterkwelder, *Schiermonnikoog*, Friesland, photographed (description incomplete, eg, regarding head pattern; photographs too blurry to determine details). **Western Black-eared Wheatear / Westelijke Blonde Tapuit** *Oenanthe hispanica* 3 October, Buytenpark, *Zoeter-*

meer, Zuid-Holland, first-winter female, photographed (@ description incomplete, especially regarding tail pattern; photographs do not show sufficient details, eg, no tail visible). **White's Thrush / Goudlijster** *Zoothera aurea* 3 October, Klompenwaard, Doornenburg, *Bemmel*, Gelderland (description incomplete); 9-10 October, Westduinpark, *Den Haag*, Zuid-Holland (description incomplete). * **Oriental White-eye / Indische Brilvogel** *Zosterops palpebrosus* 31 October, *Rijnsburg*, Zuid-Holland, photographed (@ description incomplete; based on the photographs, although clearly a *Zosterops* species, it is impossible to identify the species). **Parrot Crossbill / Grote Kruisbek** *Loxia pytyopsittacus* 13 April, Aekingerzand, *Ooststellingwerf*, Friesland, male and female (@ no description; van Manen 2004); 14-22 April, Appelscha, *Ooststellingwerf*, Friesland, male and female with three eggs, photographed (no description of the birds; egg length and width also fit Common Crossbill *L curvirostra*; photograph not conclusive; van Manen 2004); 13 May, Smilde, *Middenveld*, Drenthe, male, female and juvenile (@ no description; van Manen 2004).

2003

African Chaffinch / Afrikaanse Vink *Fringilla coelebs africana/spodiogenys* 28 November to 6 January 2004, Onnerweg, *Haren*, Groningen, male, winter plumage, photographed, videoed (Brinkhuizen et al 2004; bird showed odd underparts for male African Chaffinch; call described as similar to nominate Common Chaffinch *F c coelebs* is wrong for African).

2002

American Wigeon / Amerikaanse Smient *Anas americana* 2 February, *Harlingen*, Friesland, two, male, photographed (@ no description; photographs too blurry to identify the birds); 29 March, Ezumakeeg, *Dongeradeel*, Friesland, male (@ no description). **Terek Sandpiper / Terekruiter** *Xenus cinereus* 2 June, Monnikeburenpolder, *Wûnseradiel*, Friesland (@ description incomplete). **House Crow / Huiskraai** *Corvus splendens* 2 April, Camperduin, *Bergen*, Noord-Holland (insufficient details of a bird flying by).

2001

Greater Shearwater / Grote Pijlstormvogel *Puffinus gravis* 26 December, Camperduin, *Bergen*, Noord-Holland (@ description incomplete). **Pallid Harrier / Steppekiekendief** *Circus macrourus* 17-18 May, Veenzijdse Polder, *Wassenaar*, Zuid-Holland, second calendar-year female, photographed, videoed (rejected after review; Montagu's Harrier *C pygargus* not safely excluded; Dutch Birding 23: 236, plate 266, 2001). **Booted Eagle / Dwergarend** *Aquila pennata* 25 May, Heesch, *Bernheze*, Noord-Brabant, pale morph (@ description incomplete).

1998

Great Shearwater / Grote Pijlstormvogel *Puffinus gravis* 12 October, Camperduin, *Bergen*, Noord-Holland (description incomplete).

1997

Collared Pratincole / Vorkstaartplevier *Glareola pratincola* 1-5 August, Workumerwaard and Doniaburen, *Wûnseradiel*, Friesland, adult (Wiegant et al 1999) (now accepted as Oriental Pratincole *G maldivarum*).

1995

Booted Eagle / Dwergarend *Aquila pennata* 3 June, Loozerheide, *Weert*, Limburg, and Boshoverheide, *Cranendonk*, Noord-Brabant, pale morph (description incomplete).

1989

American Sandwich Tern / Amerikaanse Grote Stern *Sterna sandvicensis acuflavida* 11 February, Brouwersdam, *Schouwen-Duiveland*, Zeeland, adult, winter plumage, photographed (photographs show European Sandwich Tern *S s sandvicensis*; cf Olsen & Larsson 1994, p 88, plate 51).

1988

Siberian Chiffchaff / Siberische Tjiftjaf *Phylloscopus collybita tristis* 20-27 November, Schelle, *Zwolle*, Overijssel (rejected after review).

1931

Black-headed Bunting / Zwartkopgors *Emberiza melanocephala* 16-17 August, De Cocksdorp, *Texel*, Noord-Holland, male (description reminiscent of Red-headed Bunting *E bruniceps* in several respects; Stichting 'Vogeltrekstation Texel' 1932).

1890

Yellow-billed x Great Northern Loon / Geelsnavelduiker x IJsduiker *Gavia adamsii x immer* 4 December, Schellingwoude, *Amsterdam*, Noord-Holland, first-winter male, skin retained at Zoological Museum, Amsterdam, Noord-Holland (the CDNA is uncertain if a conclusion can be reached based on relatively few specimens of both presumed parent species; cf Roselaar et al 2006a).

Acknowledgements

The CDNA wants to thank Jo van der Coelen for correcting a mistake from a previous report. Dick Forsman, Hannu Jannes, Marc Herremans, Magnus Robb and Lars Svensson are thanked for their comments on some records. Enno Ebels took care of the final editing. Roland thanks the Infadels for their inspiration during many nightly hours, while Jan was inspired by Bloc Party during (mostly) late evening hours. Bram Rijksen kindly prepared the drawing of the Killdeer on p 345.

References

Alblas, E J 2005. DB Actueel: Sperweruil bij Westerbork. Dutch Birding 27: 449-450.

van den Berg, A B & Bosman, C A W 1999, 2001. Zeldzame vogels van Nederland – Rare birds of the Netherlands. First, second edition. Haarlem.

van den Berg, A B, van Loon, A J & McGeehan, A 2000. Aalscholver met kenmerken van Grote Aalscholver te Heel

in februari 2000. Dutch Birding 22: 21-25.
van den Bergh, L M J, Gerritse, W G, Hekking, W H A, Keij, P G M J & Kuyk, F 1979. Vogels van de Grote Rivieren. Utrecht.
van Beusekom, R F J 2001. Broedgeval van Noordse Nachtegaal in Flevoland in 1995. Dutch Birding 21: 203-208.
van Beusekom, R F J 2005. DB Actueel: Keizerarend bij Kamperhoek. Dutch Birding 27: 232-233.
Boon, L J R & Slaterus, R 2006. Cursorius jaaroverzicht 2005. Zandvoort.
Bosma, J G, de Bruin, A & Ebels, E B 2005. Haakbekken in Noord-Nederland in november 2004. Dutch Birding 27: 315-328.
Brinkhuizen, D, Heikamp, A & van den Berg, A B 2004. DB Actueel: Afrikaanse Vink in Haren. Dutch Birding 26: 84-85.
de Bruin, B & van der Vliet, R 2005. Recente CDNA-besluiten. Dutch Birding 27: 339.
Bunskoek, M 2005. DB Actueel: Kumliens Meeuw op Terschelling. Dutch Birding 27: 152-154.
Constantine, M & The Sound Approach 2006. The Sound Approach to birding: a guide to understanding bird sound. Poole.
van Dongen, R M, Haas, K & de Rouw, P W W 2005. Recente meldingen Nederland: januari-oktober 2005. Dutch Birding 27: 144-150, 220-227, 282-288, 351-358, 426-439.
van Dongen, R M, Haas, K & de Rouw, P W W 2006. Recente meldingen Nederland: november-december 2005. Dutch Birding 28: 55-64.
Driessens, G & Svensson, L 2005. Identification of Collared Pratincole and Oriental Pratincole – a critical review of characters. Dutch Birding 27: 1-35.
Ebels, E B 2003. Broedgeval van Grauwe Fitis op Schiermonnikoog in mei-juli 2003. Dutch Birding 25: 304-311.
Ebels, E B 2005a. DB Actueel: Roodkeelstrandloper en Grijze Strandloper in Wagejot. Dutch Birding 27: 364-366.
Ebels, E B 2005b. Monniksgier van Spaans-Franse herkomst in Nederland in maart-augustus 2005. Dutch Birding 27: 329-333.
Ebels, E B & Halff, R 2003. DB Actueel: Groenlandse Witstuitbarmsijs bij Huisduinen. Dutch Birding 25: 439-440.
Hovinga, R 2006. Kalanderleeuwerik in Eemshaven in mei 2005. Dutch Birding 28: 230-231.
Hovinga, R & Olthoff, M P 2005. DB Actueel: Vluchtige dwaalgasten in Eemshaven: Kalanderleeuwerik en Woestijnvink. Dutch Birding 27: 293-294.
van IJzendoorn, E J, van der Laan, J & CDNA 1996. Herziening van de Nederlandse Avifaunistische Lijst 1800-1979: tweede fase. Dutch Birding 18: 157-202.
de Jong, J 2004a. Erstfeststellung eines Sperlingskauzes (*Claucidium* [sic!] *passerinum*) in den Niederlanden. Eulen-Rundblick 51/52: 40-41.
de Jong, J 2004b. Dwerguil (Dwerchûle), nieuwe soort voor Nederland. Vanellus 57: 50-51.
van der Leij, R-J & Ebels, E B 2005. DB Actueel: Killdeerplevier bij Rottige Meente. Dutch Birding 27: 233-234.
van der Leij, R-J & Ebels, E B 2006. Killdeerplevier bij Rottige Meente in april 2005. Dutch Birding 28: 7-10.
Lensink, R 1993. Vogels in het hart van Gelderland. Utrecht.
Luiten, T 2005. DB Actueel: Schreeuwarend op Walcheren. Dutch Birding 27: 444-445.
van Manen, W 2004. Broedgevallen van Grote Kruisbek *Loxia pytyopsittacus* in het Drents-Friese Wold in 2004. Drentse Vogels 18: 49-53.
Meininger, P L 2005. DB Actueel: Roodoogvireo te Westkapelle. Dutch Birding 27: 445-446.
Newson, S E, Hughes, B, Russell, I C, Ekins, G R & Sellers, R M 2004. Sub-specific differentiation and distribution of Great Cormorants *Phalacrocorax carbo* in Europe. Ardea 92: 3-10.
Nieuwstraten, E & Zekhuis, M 1997. DB Actueel: Oosterse Vorkstaartplevier in Workumerwaard en bij Doniaburen. Dutch Birding 19: 211-212.
Olsen, K M & Larsson, H 1994. Sterns van Europa en Noord-Amerika. Haarlem.
Olthoff, M P 2006. Woestijnvink in Eemhaven in juni 2005. Dutch Birding 28: 235-236.
Pennington, M & Maher, M 2005. Greenland, Iceland & Hornemann's Redpolls in Britain. Birding World 18: 66-78.
Plomp, M, Berlijn, M, Alards, H, Bernardus, S, Bot, M, Ebels, E B, Edelaar, L, Levering, H, van der Maat, G, Menkveld, E, Opperman, E, Sponselee, R & Wight, H-J 2006. Dutch Birding jaaroverzicht 2005. Dvd. Linschoten.
Reijnders, C & Ebels, E B 2004. Grote Tafeleend bij Castricum aan Zee in januari-maart 2003. Dutch Birding 26: 180-184.
Rogers, M J & the Rarities Committee 2004, 2005. Report on rare birds in Great Britain in 2003; in 2004. Br Birds 97: 558-625; 98: 628-694.
Roselaar, C S, Prins, T G, Aliabadian, M & Nijman, V 2006a. Hybrids in divers (Gaviiformes). J Ornithol 147: 24-30.
Roselaar, C S, van den Berg, A B, van Loon, A J & Maassen, E 2006b. Siberische Sprinkhaanzangers in Noord-Holland in september 2002-05. Dutch Birding 28: 273-283.
Rijpma, U & Bakker, A 2006. Perzische Roodborst bij Pietersbierum in oktober 2005. Dutch Birding 28: 302-304.
van Rijswijk, W 2004. Woestijnvink op Maasvlakte in mei 2003. Dutch Birding 26: 185-189.
Stichting 'Vogeltrekstation Texel' 1932. Eerste jaarverslag der stichting en Mededeelingen betreffende de waarnemingsperioden herfst 1931 en voorjaar 1932.
van der Vliet, R E, van der Laan & CDNA 2001, 2004, 2005. Rare birds in the Netherlands in 2000; in 2003; in 2004. Dutch Birding 23: 315-347; 26: 359-384; 27: 367-394.
Wassink, A & Ebels, E B 2005. DB Actueel: Waarschijnlijke Oosterse Gekraagde Roodstaart op Texel. Dutch Birding 27: 366, III.
van Wetter, P, Alblas, E J & Ottens, G 2006. Sperweruil bij Westerbork in oktober 2005. Dutch Birding 28: 292-296.
Wiegant, W M, Steinhaus, G H & CDNA 1996. Rare birds in the Netherlands in 1994. Dutch Birding 18: 105-121.
Wiegant, W M, de Bruin, A & CDNA 1999. Rare birds in the Netherlands in 1998. Dutch Birding 21: 309-329.
Winters, R 2004. DB Actueel: Marmereend bij Pannerden. Dutch Birding 26: 357-358.
Wolf, P A 2003. DB Actueel: Orpheusgrasmus op bedrijventerrein in Middelburg. Dutch Birding 25: 443-444.

Roland E van der Vliet, Adenauerlaan 113, 3527 RC Utrecht, Netherlands
Jan van der Laan, Brouwerstraat 19, 1814 HX Alkmaar, Netherlands
Max Berlijn, Wilhelminastraat 9, 6285 AS Epen, Netherlands
CDNA, Postbus 116, 2080 AC Santpoort-Zuid, Netherlands (cdna@dutchbirding.nl)

Masters of Mystery

SWAROVSKI
OPTIK

Solutions of fifth round 2006

The solutions of the mystery photographs IX and X (Dutch Birding 28: 315, 2006) of the 2006 Masters of Mystery competition appear below.

IX Although the size of a bird is generally difficult to assess in a single photograph, the mystery bird seems to be one of the smaller gulls *Larus*. This is further supported by the thin leg and the pale coloration of the underside of the outer primary. In immature large gulls, the outer primary is dark brownish, while in adults this is either all-white or largely black.

The mystery bird is a small gull species; either one of the 'two-year gulls' or 'three-year gulls' (passing through two or three complete moult cycles before attaining adult plumage, respectively). The presence of a dark tail-band shows that these feathers are either of the first or the second generation. As second-generation tail-feathers (and older ones) are clean white or show an irregularly shaped tail-band, the evenly broad, black tail-band leads us to a bird with first generation tail-feathers. Considering that the photograph was taken in July, and as all visible feathers (for instance the underwing-coverts, primaries and tail-feathers) seem to be fresh and undamaged, the bird must be in its first calendar-year.

The underside of the outer primary of the mystery bird is largely pale and the tail has an ordinary shape: it is neither forked nor triangular. Furthermore, the black tail-band is evenly broad across t1-5, excluding Ross's Gull *Rhodostethia rosea* and Sabine's Gull *L sabini*. Of the small gulls, only Little *L minutus*, Slender-billed *L genei*, Bonaparte's *L philadelphia*, Black-headed *L ridibundus*, Relict *L relictus* and Mediterranean Gull *L melanocephalus* show these features. Slender-billed, Bonaparte's and Black-headed all show an outer primary with a long clean white tongue on the underside, strongly contrasting with the sharply defined black outer web, inner web (except in Bonaparte's) and tip, rather different from the pattern shown by the mystery bird, which shows a largely grey outer web converging into black at the tip and a clean white patch at the base. In addition, both Slender-billed and Bonaparte's show paler legs than the mystery bird. Another argument for excluding these species (as well as Little Gull) are the large grey tips to the greater under primary-coverts. This feature only fits Relict and Mediterranean. Note also the dark markings along the wing-bend. Relict can be excluded, as this species would show a small white spot near the tip of the inner web, which is absent in the mystery bird. This leaves us with Mediterranean as the correct answer.

This juvenile Mediterranean Gull was photographed by Chris van Rijswijk on 30 July 2006 at Sainte Anne-la-Palud, Finistère, France. Another photograph of the same bird is shown in plate 513. It was correctly identified by only 14% of the entrants, while most of the remaining entrants voted for Little (36%) and Bonaparte's Gull (18%).

513 Mediterranean Gull / Zwartkopmeeuw *Larus melanocephalus*, juvenile, Sainte Anne-la-Palud, Finistère, France, 30 July 2006 *(Chris van Rijswijk)*.

X Most entrants considered the large raptor of mystery photograph X to be one of the larger vultures or a species of fish eagle *Haliaeetus*. The mystery bird is characterized by a rather uniform dark brown plumage. The tail is short and the wings are well-fingered and rather broad. No signs of active moult are visible and all flight-feathers seem to be undamaged, creating a uniform and smooth trailing edge to the wing and indicating that the bird is a fresh juvenile. The bill is just visible and is grey with a dark blackish tip. Furthermore, the bill is rather small and this excludes easily all larger vultures because these all show strong and quite long bills. One of the most mentioned incorrect answers was Eurasian Black Vulture *Aegypius monachus* which has a massive black bill with greyish bill-base. In addition, the upperparts in Eurasian Black are very dark brown-black, especially in juvenile plumage, and usually do not show pale tips to the greater upperwing-coverts as shown in the mystery bird. Another difference can be found in the wing formula. In Eurasian Black, the fourth primary (p4) is long (as are p5-p10), creating an extra 'finger' on the wingtip, whereas in the mystery bird p4 is rather short. The rather uniform blackish-brown appearance of the mystery bird, prompted some entrants to vote for Hooded Vulture *Necrosyrtes monachus*. Note, however, that this species has a long and slender bill and a pale pinkish-grey (in juveniles) facial skin and throat. Both characters are absent in the mystery bird and Hooded Vulture can, therefore, be excluded.

[*Dutch Birding 28: 366-368, 2006*]

514-515 Bateleur / Bateleur *Terathopius ecaudatus*, juvenile, Kalahari Gemsbok National Park, North Cape, South Africa, 12 March 1996 *(Arnoud B van den Berg)*

The mystery bird also does not fit one of the *Haliaeetus* species, of which Bald Eagle *H leucocephalus* was mentioned most. Again, the bill of the mystery bird is much too small to fit one of the fish eagles. Furthermore, all fish eagles show a rather strongly patterned upperwing in juvenile plumage with pale or brownish tips to the greater and median coverts, a pale mantle and in most cases a whitish uppertail, creating a strong variegated pattern, quite unlike the rather uniform upperwing of the mystery bird.

If the bird is not one of the fish eagles and also not one of the larger vultures, then what can it be? The answer is found in the very short tail and the pale coloration of the inner primaries. Both are characteristic features of juvenile Bateleur *Terathopius ecaudatus*. This Bateleur was photographed by Arnoud van den Berg on 12 March 1996 in Kalahari Gemsbok National Park, North Cape, South Africa. Because the outer primaries are spread, the pointed wing so characteristic for this species is not visible. Two other photographs from the same locality, one showing the typical pinched wing-tips and strongly bulging secondaries, are shown in plate 514-515. Bateleur is a common and widespread raptor in Africa. It is on the Western Palearctic list based on a few records in the Middle East (Egypt, Iraq and Israel) and one in Tunisia. It was correctly identified by 36% of the entrants. The most mentioned incorrect answers were Eurasian Black Vulture (31%) and a wide variety of other raptors, of which three species of fish eagle (in total 15%) were most frequent.

In the fifth round for 2006, there were 39 entrants of which only six managed to identify both mystery birds correctly. From them, Martin Kühn (Germany) was drawn as the winner of a copy of the MP3 sound collection *Birds of Mexico* by Peter Boesman, donated by Birdsounds.nl. After five rounds, the number of leading

Mystery photograph XI (September)

Mystery photograph XII (January)

entrants was further reduced to two, both of them with 10 correct identifications, Julia Bayer (Germany) and Tommy Holmgren (Sweden). Hot on their heels are Chris Batty (England), Martin Kühn and Clemens Portofée (Germany) with nine correct answers. They are closely followed by four entrants with eight and 10 with seven correct identifications. The names of all entrants with at least one correct identification can be viewed at www.dutchbirding.nl.

Sixth round 2006

Photographs XI and XII represent the final sixth round of the 2006 competition. Please, study the rules (Dutch Birding 28: 37, 2006) carefully and identify the birds in the photographs. Solutions can be sent in three different ways:

- by *postcard* to Dutch Birding Association, Postbus 75611, 1070 AP Amsterdam, Netherlands
- by e-mail to masters@dutchbirding.nl
- from the website of the Dutch Birding Association at www.dutchbirding.nl

Entries for the sixth round have to arrive by **1 January 2007**. Please, indicate if you are subscribed to Dutch Birding. From those entrants having identified both mystery birds correctly, one person will be drawn who will receive a copy of *Birds of Southern Africa* by Ian Sinclair, Phil Hockey and Warwick Tarboton. Swarovski Benelux has generously agreed to sponsor this competition again in 2006. This year, the overall winner after six rounds will receive a pair of marvelous 10x32 EL binoculars.

Rob S A van Bemmelen, Stavangerweg 535, 1013 AX Amsterdam, Netherlands (masters@dutchbirding.nl)
Dick Groenendijk, Elzenstraat 14, 4043 PB Opheusden, Netherlands (masters@dutchbirding.nl)

CDNA-mededelingen

CDNA-postadres gewijzigd Vanwege het toenemend gebruik van internet is besloten de aparte CDNA-postbus (Postbus 45, 2080 AA Santpoort-Zuid) te laten vervallen en voortaan gebruik te maken van het postbusnummer dat ook door de redactie van Dutch Birding wordt gebruikt. Het nieuwe adres is: CDNA, Postbus 116, 2080 AC Santpoort-Zuid, Nederland. LAURENS STEIJN

Aankondigingen & verzoeken

Ornithologisch jaarverslag Texel 2005 In augustus 2006 verscheen het *Ornithologisch jaarverslag Texel 2005*, uitgegeven door de Vogelwerkgroep Texel. In dit negende verslag van de vogelwerkgroep wordt een systematisch overzicht gegeven van de vogelwaarnemingen op Texel, Noord-Holland, in 2005. Het boekje omvat 56 pagina's en is geïllustreerd met diverse kleurenfoto's van op Texel waargenomen vogels. Het is in een aantal boekhandels op Texel en bij EcoMare en de VVV Texel verkrijgbaar voor EUR 6.50 maar is ook te bestellen door EUR 8.00 (incl verzendkosten) over te maken op girorekening 5312854 tnv Vogelwerkgroep Texel te Den Burg ovv 'Jaarverslag 2005' en naam en adres.

Geïnteresseerden kunnen de meest recente waarnemingen lezen op de website van de Vogelwerkgroep Texel (www.vogelwerkgroeptexel.nl). Tevens kunnen via deze website eigen waarnemingen op Texel worden doorgegeven.

Corrigendum

In het bijschrift bij plaat 440 (Dutch Birding 28: 320, 2006) werd helaas niet de juiste fotograaf vermeld. De foto werd gemaakt door János Oláh. REDACTIE

In the caption of plate 440 (Dutch Birding 28: 320, 2006) unfortunately the wrong photographer was mentioned. The photograph was taken by János Oláh. EDITORS

DBA-nieuws

Betaling van abonnementsgeld voor 2007 Bij dit nummer van Dutch Birding treft u een rekening aan voor het abonnementsgeld voor 2007. Nederlandse abonnees kunnen het abonnementsgeld betalen met de eveneens bijgevoegde acceptgirokaart. Wij verzoeken de Belgische abonnees om de betaalinstructies op de rekening te volgen. In verband met gestegen productiekosten en de verhoging van de portokosten met ingang van 2007 hebben wij de abonnementsprijs iets moeten verhogen. De abonnementsprijs wordt EUR 34.00 voor Nederland en EUR 37.00 voor België. Wij verzoeken u het abonnementsgeld voor 1 januari 2007 over te maken. In verband met de kosten die verbonden zijn aan het versturen van herinneringen en aanmaningen zal voor latere betalers EUR 2.00 extra in rekening worden gebracht.

Voorts verzoeken wij u om alle correspondentie over abonnementszaken te richten aan: Dutch Birding Association, p/a Jeannette Admiraal, Iepenlaan 11, 1901 ST Castricum, Nederland, of aan circulation@dutchbirding.nl.

Payment of subscription for 2007 Enclosed with this issue, you will find the invoice for the subscription fee for 2007. The rates for 2007 will be EUR 37.00 for subscribers in Europe and EUR 40.00 for subscribers outside Europe. We kindly request to follow the instructions on the enclosed invoice. Subscribers who have not paid by 1 January 2007 will be charged EUR 2.00 extra to compensate administration costs.

For all subscription matters, please use the following address: Dutch Birding Association, c/o Jeannette Admiraal, Iepenlaan 11, 1901 ST Castricum, Netherlands, or circulation@dutchbirding.nl.

Uitverkochte nummers op DBA-website en invoering digitaal abonnement De jaargangen 1-12 van Dutch Birding zijn al geruime tijd geheel of grotendeels uitverkocht. Dit betekent dat mensen die in de jaren daarna abonnee zijn geworden vaak geen complete set meer kunnen krijgen en dat de informatie die in deze nummers is gepubliceerd niet meer optimaal toegankelijk is. Om dit probleem te verhelpen zijn deze jaargangen gescand en in pdf-formaat omgezet. Via de website van de DBA (www.dutchbirding.nl/journal/backpdf.php) zijn de nummers in te zien en te downloaden. Hieraan zijn geen kosten verbonden. De kwaliteit van de gescande nummers is niet te vergelijken met de hoge kwaliteit van het oorspronkelijke drukwerk maar voldoende om de teksten te lezen en foto's te bekijken. Redactie en bestuur hopen dat met deze service in een behoefte wordt voorzien en dat de mogelijkheid om kennis te nemen van oude artikelen en daarnaar te verwijzen volop benut zal worden. De jaargangen 13-26 zullen binnenkort (waarschijnlijk rond de jaarwisseling) als 'back issue' in pdf-formaat worden aangeboden; tegen betaling van een vergoeding (50% van de kosten voor een gedrukt exemplaar) zijn deze pdf-bestanden dan via de DBA-website te bestellen. Later in 2007 zal ook jaargang 27 als pdf-bestand beschikbaar komen. Van jaargang 16-26 (en natuurlijk 27-28) zijn momenteel ook nog voldoende gedrukte exemplaren beschikbaar.

Met ingang van 2007 maken redactie en bestuur een begin met het aanbieden van een digitaal abonnement ('e-abonnement'). Hierbij wordt het nummer in pdf-formaat per e-mail verzonden tegen een aanmerkelijk lager tarief dan bij een normaal abonnement. Deze service wordt in eerste instantie aangeboden aan abonnees buiten Europa omdat de hoge portokosten en lange verzendtijd per (zee-)post voor deze groep het meest nadelig zijn. Bij voldoende belangstelling en goed functioneren zal het digitale abonnement met ingang van 2008 ook voor abonnees in Europa worden ingevoerd.

Sold-out back issues on DBA-website and introduction of digital subscription Volumes 1-12 of Dutch Birding have been sold out completely or largey. The issues of volume 1-12 have been scanned and can now be viewed in pdf-format on the website of the Dutch Birding Association (www.dutchbirding.nl/journal/backpdf.php) and can be downloaded free of charge. The quality of the scans does not match the printed version but the scans give access to both texts and photographs and enable reference to publications that would otherwise receive less attention. As a next step, issues of volumes 13-15 and 16-26 (of which hard copy back issues are still available) can soon be ordered as pdf for a small fee (50% of the price of a printed issue). It is expected that these electronic back issues will be available with the start of the new year. Pdf files of volume 27 will follow later in 2007.

Starting in 2007, a digital subscription ('e-subscription') will be introduced, with subscribers receiving a pdf of each new issue by email. The digital subscription is introduced first for subscribers outside Europe only (because postage costs and long delivery periods are most disadvantageous for this group). The subscription rate is considerably lower than a regular subscription (overseas subscription rates for 2007: hard copy EUR 40.00, digital copy EUR 20.00, combined EUR 55.00). In 2008, introduction of the digital subscription is planned also for subscribers in Europe.

Geanimeerde Texelweek 2006 Toen op zaterdag 7 oktober 2006 de Dutch Birding-vogelweek op Texel, Noord-Holland, begon was de zeetrek erg goed, met het tegenwoordig gebruikelijke palet aan interessante meeuwen, jagers en pijlstormvogels, inclusief Vale Pijlstormvogel *Puffinus mauretanicus*. Een Pallas' Boszanger *Phylloscopus proregulus* bij Dorpszicht was slechts voor een enkeling weggelegd. Er werden minimaal twee Bladkoningen *P inornatus* gemeld. Zondag

zorgden een overvliegende Zwarte Ibis *Plegadis falcinellus* en Grote Pieper *Anthus richardi* en een zingende Europese Kanarie *Serinus serinus* in Den Burg voor enige rimpelingen in het vlakke beeld van dit weekend. Jammer genoeg laat in de middag werd aan de Jan Ayenslag een Aziatische Roodborsttapuit *Saxicola maurus* ontdekt, die slechts door een handvol vogelaars nog getwitcht kon worden.

Maandag was een Zwarte Zeekoet *Cepphus grylle* op zee bij Ecomare het zoveelste bewijs dat deze fraaie zeevogel een steeds vertrouwdere verschijning wordt in onze noordelijke kustwateren. De gemelde Pallas' Boszanger in De Cocksdorp was een tweemanswaarneming. Dinsdag ontstond er dan eindelijk enige echte beroering. Wim Wiegant ontdekte in de haven van Oudeschild namelijk een spreeuw *Sturnus* die wel érg veel op een Zwarte Spreeuw *S unicolor* in vol zomerkleed leek. Alle mogelijke en onmogelijk semafooncodes werden in de soms verhitte strijd geworpen om de discussie over de vogel gaande te houden. Helaas zal de op zich interessante vogel de boeken ingaan als een hormoongestoorde Spreeuw *S vulgaris*. Deze vogel trok zoveel aandacht dat er drie dagen lang nauwelijks iets anders werd gemeld van het eiland. Op donderdag 12 oktober zorgden twee Zwarte Rotganzen *Branta nigricans* voor enige afwisseling. De claim van een Vale Lijster *Turdus obscurus* in een tuin bleek helaas wel erg vaag, en een Siberische Tjiftjaf *P collybita tristis* op de noordpunt van het eiland zal nog in de slag moeten gaan met de CDNA. Dat de Texelweek officieel was afgesloten weerhield een Kleine Vliegenvanger *Ficedula parva* er niet van zich in het weekend van 14 en 15 oktober uitgebreid te laten bewonderen in een Texelse achtertuin.

Ondanks de afwezigheid van echte toppers werd de week behalve door de 'zwarte' Spreeuw levendig gehouden door het lezingenprogramma en de Big Day. Van de internetverbinding in het gezellige hoofdkwartier De Robbenjager (waar op zaterdag en donderdag Swarovski een stand had) werd druk gebruik gemaakt. Vrijwel elke dag bleken er zich toch c 100 gelouterde vogelaars op het eiland te bevinden. Een groot deel van hen kon op de zaterdagavond genieten van de vertrouwde kwaliteit van de opnamen van Chris Schenk (Hokkaido, Japan, in de winter). Speciale aandacht was er deze avond voor Texel-veteraan Wim Wiegant. Aan maar liefst 25 volledige DB-vogelweken heeft hij meegedaan (vier op Terschelling, waaronder enkele dubbele, en 21 op Texel). Wim is er in al die jaren maar één dag tussenuit geweest vanwege een verplichting aan de wal. Dat kostte hem de waarneming van een Geelsnavelkoekoek *Coccyzus americanus* (uiteindelijk niet aanvaard...), nota bene in de tuin van zijn vakantiehuisje. Theo Admiraal maakte zondag zijn debuut als spreker met een zeer informatief reisverslag van zijn trip met hindernissen door Venezuela. Texelaar en wereldreiziger Arend Wassink maakte zich maandagavond verdienstelijk met een ingelaste lezing over Kazachstan en Antarctica. Dick Groenendijk had zich weer ingespannen voor een goed verzorgde mystery bird-competitie op de dinsdagavond. De drie winnaars (Jan Kees Bosschenbroek, Kees de Mooij en Edwin de Weerd) konden kiezen tussen *Threatened birds of the world, Currassows and related birds* (beide fraaie boeken beschikbaar gesteld door uitgeverij Lynx Ediciones) of een dvd naar keuze van Plomp Digital Video. Chris van Rijswijk is een tamelijk nieuwe ster aan het vogelfotografenfirmament en gaf op woensdag een staalkaart van zijn kunnen, met veel Nederlandse vogels én veel vogelaars in actie. Diederik Kok schroomde niet enkele intieme details van het Afrikaanse dierenleven te laten zien in zijn ontspannen verslag van een familie-uitje met veel vogels in Tanzania.

Aan de Big Day op donderdag 12 oktober deden zeven teams mee. Dit jaar lukte het het vierkoppige team van Dick Groenendijk dan eindelijk om de felbegeerde Swarovski-rugzakken 'type Shetland' te winnen, beschikbaar gesteld door onze gewaardeerde sponsor Swarovski. De score was 118 soorten. Het team van Marco Witte werd verblijd met Swarovski-t-shirts en voor alle deelnemers aan de wedstrijd was er een door Grasduinen geschonken agenda. De uitslag van de 'Soort van de Week' was op dat moment helaas niet zo eenduidig. Swarovski heeft daarvoor een nieuw carbonstatief ter waarde van EUR 580 beschikbaar gesteld. Tot verdriet van Wim Wiegant heeft zijn 'Zwarte Spreeuw' het niet gehaald. Op de aankomende Dutch Birding-vogeldag (17 maart 2007) wordt de prijs naar verwachting uitgereikt. Vooralsnog maakt de ontdekker van de Aziatische Roodborsttapuit de meeste kans. Gijsbert van der Bent, Wietze Janse & Marc Plomp

Dutch Birding-vogelweek op 13-18 oktober 2007 Voor volgend jaar staat de Dutch Birding-vogelweek op Texel, Noord-Holland, gepland van zaterdag 13 oktober tot en met donderdag 18 oktober 2007. Een deel van Nederland heeft dan herfstvakantie, waardoor onder andere meer scholieren en studenten de kans krijgen deel te nemen. Het jaar daarna (2008) zal de vogelweek opnieuw plaatsvinden in de herfstvakantie, maar dan in de vakantieperiode van een ander deel van Nederland. De vogels hebben we niet aan een touwtje maar samen met sponsor Swarovski beraadt de organisatie zich op manieren om nog meer vogelaars naar het eiland te lokken en bij de waarnemingen te betrekken. Er komt in ieder geval een logboek op een vaste plek. Ook zal er iedere avond een nabespreking zijn van alles wat er die dag gezien is. Over het programma zal men in de zomer van 2007 worden geïnformeerd. Gijsbert van der Bent, Wietze Janse & Marc Plomp

WP reports

This review lists rare and interesting birds reported in the Western Palearctic mainly in **late September-early November 2006**. The reports are largely unchecked and their publication here does not imply future acceptance by a rarities committee. Observers are requested to submit their records to each country's rarities committee. Corrections are welcome and will be published.

GEESE TO DUCKS Rare geese in the Azores included three **Pink-footed Geese** *Anser brachyrhynchus* at Albarnaz lighthouse, Flores, on 9 October and one at Achada das Furnas, São Miguel, on 22-24 October, one **Barnacle Goose** *Branta leucopsis* at Ribeira do Mainho, Ponta Delgada, Flores, from 30 October to 5 November, and **Pale-bellied Brent Geese** *B hrota* (probably of Nearctic origin) at Ponta Delgada on 3 October and 3 November, at Cabo da Praia, Terceira, from 9 October to 2 November, at Santa Cruz, Flores, on 25 October and at Fajã Grande, Flores, on 28 October; there were only one or two previous records for these three species. In the Netherlands, the number of **Red-crested Pochards** *Netta rufina* at a small pond at Harderbroek, Flevoland, increased to an unprecedented 375 on 26 October. On 30 September, the male **Canvasback** *Aythya valisineria* at Castricum, Noord-Holland, the Netherlands, returned for its 'fifth' winter; it was first seen on 9 January 2003 but was not noticed in the winters of 2003/04 and 2004/05. A male **Boreal Eider** *Somateria mollissima borealis* stayed on Fair Isle, Shetland, Scotland, from 5 October onwards. The male **Black Scoter** *Melanitta americana* first seen at Llanfairfechan, North Wales, on 10 March 1999 returned on 24 September for its ninth winter. In the Netherlands, one or two adult male **Buffleheads** *Bucephala albeola* were swimming again at various sites along the IJsselmeer coast of Noord-Holland from 17 September to 4 October; from 18 November, this or another male was back at Barendrecht, Zuid-Holland, where it also stayed during the previous two winters. A male on Unst on 12-14 November was the first for Shetland. The male **Barrow's Goldeneye** *B islandica* first present at Quoile Pondage, Down, Northern Ireland, from 20 November 2005 to 8 April returned on 4 November. Another male was found at Callander, Forth, Scotland, on 19 November. The male **American Black Duck** *Anas rubripes* at Garður, Iceland, remained throughout the period and the one in A Coruña, Spain, until at least 5 October. Another was seen at Ventry Beach, Kerry, Ireland, from 20 to at least 23 October. The first for Fair Isle was present from 1 November onwards. An isotope analysis of the feathers of a first-year female **Baikal Teal** *A formosa* shot at Skælskør, Sjælland,

516 Great Blue Heron / Amerikaanse Blauwe Reiger *Ardea herodias*, Fajã Granda, Flores, Azores, 31 October 2006 *(Kris De Rouck)*

Denmark, on 24 November 2005 (Dutch Birding 28: 38, 2006) in good condition, which had started to moult the juvenile feathers, made clear that it was a genuine vagrant; the juvenile feathers showed values as in the species' breeding areas in Siberia, while those of the new feathers were similar to values from Mallards *A platyrhynchos* in western Europe. In September, several were found in the western Aleutians, Alaska, USA. In England, three single **Marbled Ducks** *Marmaronetta angustirostris* were at Stanpit Marsh, Dorset, from 23 September to 16 October (juvenile), at Rye Harbour, East Sussex, from 25 September to 29 October and at North Warren, Suffolk, on 28 October.

LOONS TO HERONS A Danish record count concerned 10 **Yellow-billed Loons** *Gavia adamsii* flying south-east past Skagen, Nordjylland, on 20-22 October. In Ireland, an adult was seen off Cape Clear Island and in Roaringwater Bay, Cork, from 26 October. An immature **Black-browed Albatross** *Thalassarche melanophris* flew past Utsira, Rogaland, on 8 October; up to 2002, there are 11 records for Norway. A **Herald Petrel** *Pterodroma arminjoniana* was seen for an hour south of Corvo, Azores, on 29 October. Intriguingly, peak breeding activity on the nearest known colonies at Trinidade and Martin Vaz islands off north-eastern Brazil is from October to April. In August, an amazing seabird discovery took place on the mainland of south-western France where three nests of **Scopoli's Shearwater** *Calonectris diomedea* were found west of Bordeaux at Bassin d'Arcachon, Gironde, constituting the first breeding records of this Mediterranean species for the Atlantic (the nearest Cory's Shearwaters *C borealis* breed 615 km to the south in the Berlengas, Portugal). It now appears that already since 30 years occasionally up to 100 individuals were observed here while behaving as if they were close to a colony (singing and copulating very close to the shore) and that in 1988 in the same area eight were tape-lured, trapped and identified as Scopoli's, but it took until 2005 before a recently fledged juvenile was caught on land. Curiously, this summer's nests were found in a suburban area in holes underneath a cottage and a high tree (Ornithos 13: 316-319, 2006). It is also noteworthy that, recently, a 'pure' colony of **Cory's Shearwater** was discovered in Almería on the Mediterranean coast of Spain (Molecular Phylogenetics and Evolution 41: 322-332, 2006). The first-ever twitchable Cory's for Denmark remained off Gilleleje, Nordsjælland, for 90 min on 8 October. A **Yelkouan Shearwater** *Puffinus yelkouan* flying past Cap Gris Nez, Pas-de-Calais, at 16:05 on 12 November was the fifth claim of this species for north-western France (the first was in 1998). If accepted, a **Madeiran Storm-petrel** *Oceanodroma castro* reported off Wangerooge, Niedersachsen, on 29 October would be the first for Germany. In Poland, at least three **Pygmy Cormorants** *Phalacrocorax pygmeus* remained at Jeziorsko and Przemkow until at least 19 October. The first **Green Heron** *Butorides virescens* for the Netherlands first present in juvenile plumage at De Nieuwe Meer, Amsterdam, Noord-Holland, from 25 April to 9 June reappeared in summer plumage at another Amsterdam site on 14 September (according to local information, it was present here since mid-July) and was last seen on 30 September (cf Dutch Birding 28: 243, 2006). The sixth **Squacco Heron** *Ardeola ralloides* for the Cape Verde Islands was found at Ribeira Brava, São Nicolau, on 27 September. A **Cattle Egret** *Bubulcus ibis* at Niemelänkylä, Ylivieska, on 5-8 October and possibly the same at Kaihuanmutka, Rovaniemi, from 9 October was (only) the second for Finland; the first occurred in May 2002. In the Netherlands, not only the number of breeding pairs of **Great Egret** *Casmerodius albus* (cf Dutch Birding 28: 325, 2006) increases steadily but also that of **Little Egret** *Egretta garzetta*, with 125 pairs of the latter in 2006 of which 55 at Quackjeswater, Zuid-Holland, and 25 in Zeeuws-Vlaanderen, Zeeland. At least 21 **Bourne's Herons** *Ardea bournei* (eight adults, three subadults and 10 juveniles) and seven 'active' nests were counted at the world's only colony of this species in a solitary tree at Liberão, Santiago, Cape Verde Islands, on 15 October. More than 3300 young **Greater Flamingos** *Phoenicopterus roseus* fledged from the new colony at the Macchiareddu saltpans, Sardinia, Italy. In Hungary, a juvenile was photographed at Rétszilas, Fejér, on 19 November.

RAPTORS TO CRANES At Tarifa, Cádiz, Spain, at least four **Rüppell's Griffon Vultures** *Gyps rueppellii* were present, including a juvenile, on a carcass at La Janda on 9 September; on 6 October, this or another bird was trapped at Algeciras and released at Algarrobo bird observatory where it was still present on 10 October. The first **Black-eared Kite** *Milvus lineatus* for Britain and western Europe at Fosdyke, Lincolnshire, was a juvenile discovered on 2 November and identified from images posted on the internet almost two weeks later when it was still present. **Long-legged Buzzards** *Buteo rufinus* were seen, for instance, at Hanko, Finland, on 14-15 September (12th record), in the Beskid Zywiecki mountains, Poland, on 17 September (adult) and in Ticino, Switzerland, on 26 September. In the Camargue, Bouches-du-Rhône, France, possibly the same individual first reported on 21 October was seen daily at Mas Neuf de Vaccarès from 5 November onwards. As usual, one or two immature **Greater Spotted Eagles** *Aquila clanga* were present in the Camargue from 28 October onwards. The first breeding pair of **Lesser Spotted Eagle** *A pomarina* for France produced one young both in 2003 and 2005 in Franche-Comté (Ornithos 12: 310-320, 2005); there are now 25 diurnal raptor species breeding in France, the rarest in 2005 being **Black-winged Kite** *Elanus caeruleus* (15 pairs), **Lammergeier** *Gypaetus barbatus* (41), **Egyptian Vulture** *Neophron percnopterus* (74), **Eurasian Black Vulture** *Aegypius monachus* (17), **Bonelli's Eagle** *A fasciata* (29), **Osprey** *Pandion haliaetus* (53) and **Red-footed Falcon** *Falco vespertinus* (1-2) (Ornithos 13: 174-191, 2006). In Sweden, a **Steppe Eagle** *A nipalensis* on Öland on 5-13 October was trapped and ringed on 10 October. Apart from two more usual **Amur Falcons** *F amurensis*, also

517 Long-billed Murrelet / Aziatische Marmeralk *Brachyramphus perdix*, Dawlish Warren, Devon, England, 11 November 2006 *(Steve Young/Birdwatch)*

518 Long-billed Murrelet / Aziatische Marmeralk *Brachyramphus perdix*, Dawlish Warren, Devon, England, 12 November 2006 *(Rebecca Nason)*

519 American Great Egret / Amerikaanse Grote Zilverreiger *Casmerodius albus egretta*, Povoação, São Miguel, Azores, 30 September 2006 *(Daniele Occhiato)* **520** Bourne's Heron / Kaapverdische Purperreiger *Ardea bournei*, Liberão, Santiago, Cape Verde Islands, 15 October 2006 *(Edwin Winkel)* **521** Boreal Eider / Noordelijke Eider *Somateria mollissima borealis*, immature male, Fair Isle, Shetland, Scotland, 8 October 2006 *(Rebecca Nason)*

522 Least Sandpiper / Kleinste Strandloper *Calidris minutilla* (left) and Semipalmated Sandpiper / Grijze Strandloper *C pusilla*, Lajes do Pico, Pico, Azores, 19 September 2006 *(Sébastien Siblet)*

523 Short-billed Dowitcher / Kleine Grijze Snip *Limnodromus griseus*, juvenile, Lajes do Pico, Pico, Azores, 22 September 2006 *(Sébastien Siblet)*

the first **Red-footed Falcon** for the United Arab Emirates was found at Fujairah national dairy farm at Dibba on 14 October. An **Eleonora's Falcon** *F eleonorae* was reported at Gottröra ka, Uppland, Sweden, on 17 September. On 21 September, a pale morph was first seen leaving Falsterbo, Skåne, Sweden, and one hour and seven minutes later at Stevns Klint, Sydsjælland, Denmark. The 15th for Sweden was reported on Vinga, Västergötland, on 1 October. On 28-31 October, an unringed **Saker Falcon** *F cherrug* roosted frequently on a pylon at Zülpich, Nordrhein-Westfalen, Germany, c 50 km east of the Dutch border near Vaals, Limburg. In Italy, immatures were photographed at Fiumicino, Roma, on 15 October and at Migliarino, Pisa, on 7 November. On 18 October, a **Cape Verde Falcon** *F madens* was seen at the highest mountain of São Nicolau. In Israel, single **African Swamp-hens** *Porphyrio madagascariensis* remained at Ma'agan Michael from August until early October and at Ein Hahoresh fish-ponds, Heffer valley, during September. In Turkey, a **Grey-headed Swamp-hen** *P poliocephalus* was seen at a new site near Igdir, Tuzluca, on 8 September. The second **Common Crane** *Grus grus* for Madeira was seen at the westernmost tip of the island on 18 October and photographed the next day.

WADERS In September, numbers of Nearctic waders in Ireland, southern England and south-western France were higher than in other years. The fifth **Semipalmated Plover** *Charadrius semipalmatus* for the Cape Verde Islands was photographed south of Tarrafal, Santiago, on 17 October. In Scotland, a **Killdeer** *C vociferus* first seen on North Uist, Outer Hebrides, on 19 October turned up on Oronsay, Argyll, on 21 October. The third **American Golden Plover** *Pluvialis dominica* for Senegal was photographed at Ziguinchor, Casamance, on 15 October. The 11th for the Cape Verde Islands was present briefly on Raso on 21 October. Up to 300 **Sociable Lapwings** *Vanellus gregarius* were counted at Manych in the Stavropol region, Russia, during 1-17 September (cf Dutch Birding 27: 349, 2005). In Germany, singles occurred at Umpferstedt, Thüringen, on 25-30 September and at Halle-Seeben, Sachsen-Anhalt, on 15 October. Others were present at Kloten, Zürich, Switzerland, on 2-8 October, in Upper Silesia, Poland, on 8 October (two singles, one staying until 29 October), at Kiskunság, Hungary, on 10-11 October (juvenile) and at Nouzilly, Indre-et-Loire, France, from 10 to at least 24 October. On 20 September, a juvenile **Western Sandpiper** *Calidris mauri* was seen briefly at Keyhaven Marshes, Hampshire, England, where a **Semipalmated Sandpiper** *C pusilla* was also present. If accepted, a **Least Sandpiper** *C minutilla* at Kirchdorf am Inn, Oberösterreich, on 17 September will be the first for Austria. In mainland Spain, an unprecedented 26 **Pectoral Sandpipers** *C melanotos* were recorded during September. Six **Jack Snipes** *Lymnocryptes minimus* on Corvo on 6 October constituted the sixth record for the Azores. The sixth **Long-billed Dowitcher** *Limnodromus scolopaceus* for Iceland occurred at Eyrarbakki on 23-20 September. An **Upland Sandpiper** *Bartramia longicauda* was found together with a **Buff-breasted Sandpiper** *Tryngites subruficollis* between Sidi Moussa and Oualidia, Morocco, on 28 September. In Shetland, an **Upland Sandpiper** was observed on Unst on 6-7 October. The fifth and sixth **Marsh Sandpiper** *Tringa stagnatilis* for the Azores were found at Cabo da Praia, Terceira, on 22 September and at Lajes do Pico, Pico, on 5 October. **Wilson's Phalaropes** *Phalaropus tricolor* were seen at Walmsley and Stithians, Cornwall, on 14-25 September, at Bridges of Ross, Clare, Ireland, on 23 September, at El Matà, Aiguamolls de l'Empordà, Girona, Spain, on 26-27 September and in Charente-Maritime, France, from 7 October.

GULLS TO AUKS A **Ross's Gull** *Rhodostethia rosea* discovered at Red Hill Marina, Salton Sea, California, USA, on 18 November was the farthest south ever for North America. In England, the long-staying **Bonaparte's Gull** *Larus philadelphia* remained in Northumberland and Cleveland during October-November. Two single adults remained in Kerry, Ireland. **Laughing Gulls** *L atricilla* were found at Jökulsárlón, Iceland, on 15 September (second-winter), at Brora, Highland, Scotland, on 8-9 October (adult), at L'Albufera de València, València, on 9 October (adult; taken into care on 2 August and released on 19 September) and at Ria do Burgo, A Coruña, Spain, on 25 October (second-winter). The second for Switzerland was an adult at Luzern on 30-31 October wearing a ring which proved that it was the same individual that occurred in the Netherlands in July-August 2000, October 2001 and April 2002, at Zwillbrocker Venn, Nordrhein-Westfalen, Germany, in April-July 2001 and April-June 2002, and at San Remo, Liguria, Italy, in January-February 2004 and from November 2004 to January 2005 (cf Dutch Birding 26: 297-301, 2004, 27: 60, 2005). If accepted, an **Atlantic Yellow-legged Gull** *L michahellis atlantis* at Carrahane Strand, Kerry, from 25 September to 12 October will be the first for Ireland (Birding World 19: 391-392, 2006). An unringed second-year **Elegant Tern** *Sterna elegans* was photographed at Banc d'Arguin, Gironde, France, on 20 September and an unringed adult at Marismas del Odiel, Huelva, Spain, on 3 October. An adult **Forster's Tern** *S forsteri* stayed at Cruisetown north of Clogher Head, Louth, Ireland, from 8 October to at least mid-November. In England, a **Brünnich's Murre** *Uria lomvia* was seen briefly, twice, off East Yorkshire in association with a strong passage of Little Auk *Alle alle* on 2 November. The first **Long-billed Murrelet** *Brachyramphus perdix* for Britain was photographed at Dawlish Warren, Devon, England, on 7 November when it was misidentified as Little Auk. It was identified correctly from images posted on a local website and, subsequently, it was searched for and seen again from 11 to at least 14 November. The species' first record for the WP concerned a first-winter found dead in a fisherman's net and first misidentified as Black Guillemot *Cepphus grylle* at Zollikon, Zürichsee, Switzerland, between 15 and 18 December 1997 (Dutch Birding 20: 311, 1998, 21: 120, 1999, Br Birds 93: 190-199, 2000, Ornithol Beob

524 Semipalmated Plover / Amerikaanse Bontbekplevier *Charadrius semipalmatus*, Tarrafal, Santiago, Cape Verde Islands, 17 October 2006 *(Edwin Winkel)* **525** American Golden Plover / Amerikaanse Goudplevier *Pluvialis dominica*, Raso, Cape Verde Islands, 21 October 2006 *(Edwin Winkel)* **526** Long-billed Dowitcher / Grote Grijze Snip *Limnodromus scolopaceus*, juvenile, Cabo da Praia, Teirceira, Azores, 16 September 2006 *(Sébastien Siblet)* **527** Upland Sandpiper / Bartrams Ruiter *Bartramia longicauda*, Unst, Shetland, Scotland, 7 October 2006 *(Hugh Harrop)* **528** Least Sandpiper / Kleinste Strandloper *Calidris minutilla*, Ponta Delgada, Flores, Azores, 13 September 2006 *(Sébastien Siblet)* **529** Spotted Sandpiper / Amerikaanse Oeverloper *Actitis macularius* (left), with Common Sandpiper / Oeverloper *A hypoleucos*, Nethertown, Wexford, Ireland, 9 September 2006 *(Paul & Andrea Kelly/irishbirdimages.com)*

530 Eurasian Collared Dove / Turkse Tortel *Streptopelia decaocto*, Cabo da Praia, Teirceira, Azores, 16 September 2006 *(Sébastien Siblet)* **531** Common Crane / Kraanvogel *Grus grus*, Madeira, 19 October 2006 *(René Pop)* **532** Isabelline Wheatear / Izabeltapuit *Oenanthe isabellina*, Carmel Head, Anglesey, Wales, 23 September 2006 *(Jeroen Reneerkens)* **533** Siberian Accentor / Bergheggenmus *Prunella montanella*, Rumeli Feneri, Istanbul, Turkey, 2 November 2006 *(Soner Bekir)* **534** Presumed Daurian Shrike / vermoedelijke Daurische Klauwier *Lanius isabellinus*, Soral, Genève, Switzerland, 7 October 2006 *(Adrian Jordi)* **535** Iberian Chiffchaff / Iberische Tjiftjaf *Phylloscopus ibericus*, Oukaimeden, High Atlas, Morocco, 12 October 2006 *(Arnoud B van den Berg)*

97: 243-248, 2000). Remarkably, there are also 10s of records of this Asian species both in western and eastern North America, south to Florida, USA (Birding 29: 460-475, 1997, Dutch Birding 24: 378, 2002).

DOVES TO SWALLOWS Two **Eurasian Collared Doves** *Streptopelia decaocto* at Cabo da Praia on 16-18 September were the first for the Azores. A first-year **Oriental Turtle Dove** *S orientalis* photographed at Fedje, Hordaland, on 11 October was the eighth for Norway. In Finland, a **Western** *S o meena* stayed at Rauma, Finland, from 5 November and an **Eastern** *S o orientalis* at Kristiinankaupunki, 160 km from Turku, from 12 November. Also on 5 November, one returned at Falköping, Västergötland, Sweden, where it was first seen from 25 December 2005 to at least 19 April. In 2006, the number of known (ie, using nesting-boxes) **Common Barn Owls** *Tyto alba* in the Netherlands rose to 2726 breeding pairs, of which 559 in Gelderland and 515 in Friesland. The first **Eurasian Scops Owl** *Otus scops* for Iceland since 1975 was photographed at Höfn on 29 October. The **Snowy Owl** *Bubo scandiacus* in the Outer Hebrides, Scotland, was again seen on Lewis on 9 November. The second-winter female in Mayo, Ireland, from 9 September was last seen on 10 October and another was in Donegal from 8 October. In Poland, a **Northern Hawk Owl** *Surnia ulula* was photographed at Biebrza marshes on 30 October. The first **Pallid Swift** *Apus pallidus* for the Netherlands was on Vlieland, Friesland, on 20-21 October. If accepted, a **Little Swift** *A affinis* seen in flight and perched on a building at IJmuiden, Noord-Holland, for 5 min on 20 November will also be the first for the Netherlands. In England, a first-winter **European Roller** *Coracias garrulus* was present in Northumberland on 8-19 October. The result of recent analyses of mtDNA in museum specimens suggests that **American Ivory-billed Woodpecker** *Campephilus (principalis) principalis* and **Cuban Ivory-billed Woodpecker** *C (p) bairdii* split a million years ago and should be treated as two (extinct) species, not subspecies (see www.birds.cornell.edu/ivory/latest/woodpeckerDNA/document_view). On Raso, a survey of **Raso Larks** *Alauda razae* by two birders on 20-23 October revealed the presence of (only) 60 individuals in more than half of the available habitat; three nests were found, only one containing an egg, and no fledglings or juveniles were seen. An influx of **Eurasian Crag Martins** *Ptyonoprogne rupestris* in Sweden included four individuals at Vaggeryd, Småland, on 20-21 October, one at Kullen, Skåne, on 20 October and three at Linköping, Östergötland, on 22-25 October; there were only two previous records. If accepted, one reported briefly at Badshot Lea, Surrey, on 22 October would be the seventh for Britain. The first influx for the Netherlands became apparent in November. It started with sightings of two by two birders on 5 November at Amsterdam and two by four birders on 7 November at Westenschouwen, Zeeland. From 14 to at least 24 November, up to two were present and well documented at Hoorn, Noord-Holland (there were a few reports from previous years for the Netherlands but these were not (yet) accepted).

PIPITS TO ACCENTORS In Shetland, **Blyth's Pipits** *Anthus godlewskii* were found near Sumburgh, Mainland, on 12 October and on Fair Isle on 22-23 October. In Scilly, England, one stayed on St Mary's and Bryher on 18-26 October. In France, one turned up on Île d'Hoëdic, Morbihan, on 31 October. The second **Olive-backed Pipit** *A hodgsoni* for Italy was trapped on Ventotene, Lazio, in mid-October. In Shetland, **Pechora Pipits** *A gustavi* were present on Fair Isle on 15-23 September, on Fetlar on 18-21 September, on Foula on 9-13 October and at Pool of Virkie, Mainland, on 14-15 October. In Norway, one was found at Grimsøya, Røst, Nordland, on 22 September. In England, one was reported at Flamborough Head, East Yorkshire, on 24 September. A DNA sample taken from an **American Buff-bellied Pipit** *A rubescens rubescens* trapped in Hälsingland in December 2005 confirmed its identity as the first for Sweden (there is also one Swedish record of Siberian Buff-bellied Pipit *A r japonicus*). A **Forest Wagtail** *Dendronanthus indicus* photographed at Al Abraq on 10 November was the first for Kuwait (and the first within the WP boundaries as defined by *Birds of the Western Palearctic*). There are quite a few previous records of this species from Oman and the UAE; from 23 October to at least 7 November, up to four were present on Masirah, Oman, of which three were ringed. The first **Blue-headed Wagtail** *Motacilla flava* for the Azores was a female at Lajes do Pico, Pico, on 24 September. The ninth **Citrine Wagtail** *M citreola* for Iceland stayed at Árbær/Mýrar on 14-18 September. The first **Siberian Accentor** *Prunella montanella* for Turkey was photographed at Rumeli Feneri, Istanbul, on 2 November.

THRUSHES A late **Thrush Nightingale** *Luscinia luscinia* stayed on Fair Isle on 25-28 October. The first **Siberian Rubythroat** *Luscinia calliope* for the Faeroes was a male at Tórshavn on 11 October. On 26-28 October, one was seen in a garden at Sunderland, Durham, England. Outside Finland, extralimital **Red-flanked Bluetails** *Tarsiger cyanurus* were found on Gambell, Alaska, USA, on 30 September, on Sylt, Schleswig-Holstein, Germany, on 10 October, on Unst on 13-16 October, at Thorpeness, Suffolk, England, on 16-23 October, at Korbeek-Lo, Vlaams-Brabant, Belgium, on 17 October (trapped), at Røvær, Haugesund, Rogaland, on 19-20 October, on Whalsay, Shetland, on 21-24 October and at Brae, Mainland, Shetland, on 23 October. The second **Whinchat** *Saxicola rubetra* for the Azores was discovered near Pico do Lourenco, Flores, on 4 November. **Isabelline Wheatears** *Oenanthe isabellina* were present at Carmel Head, Anglesey, Wales, on 23-25 September, on Sein, Finistère, France, on 14-15 October, at Filey, North Yorkshire, England, on 14 October, and on Helgoland, Schleswig-Holstein, on 15 October. The final tally of a remarkable influx of **Northern Wheatears** *O oenanthe* from 8 August to 23 October in eastern North America was 41 individuals, with 12 in Nova Scotia, Canada. A **Pied Wheatear** *O pleschanka* was seen on Bryher, Scilly, on 20-21 October and two singles were found in Sweden on

536 Western Oriental Turtle Dove / Oosterse Tortel *Streptopelia orientalis meena*, juvenile, Rauma, Finland, 9 November 2006 *(Kari Eischer)*

537 Isabelline Wheatear / Izabeltapuit *Oenanthe isabellina*, Helgoland, Schleswig-Holstein, Germany, 15 October 2006 *(Axel Halley)*

538 Eurasian Crag Martins / Rotszwaluwen *Ptyonoprogne rupestris*, Vaggeryd, Småland, Sweden, 20 October 2006 *(Kent & Oscar Jönsson)*

539 Blyth's Pipit / Mongoolse Pieper *Anthus godlewskii*, Sumburgh Head, Mainland, Shetland, Scotland, 12 October *2006 (Hugh Harrop)*

540 Swainson's Thrush / Dwerglijster *Catharus ustulatus*, Ouessant, Finistère, France, 3 October 2006 *(Vincent Legrand)*

541 Hermit Thrush / Heremietlijster *Catharus guttatus*, Cape Clear Island, Cork, Ireland, 20 October 2006 *(Richard Bonser)*

542 Masked Shrike / Maskerklauwier *Lanius nubicus*, adult, Utsira, Rogaland, Norway, 3 October 2006 *(Christian Tiller)*

543 Red-flanked Bluetail / Blauwstaart *Tarsiger cyanurus*, first-winter, Thorpeness, Suffolk, England, 23 October 2006 *(Dave Stewart)*

544 Black-headed Bunting / Zwartkopgors *Emberiza melanocephala*, Toab, Shetland, Scotland, 16 September 2006 *(Hugh Harrop)* **545** Pechora Pipit / Petsjorapieper *Anthus gustavi*, Fair Isle, Shetland, Scotland, 23 September 2006 *(Paul Baxter)* **546** Grey-necked Bunting / Steenortolaan *Emberiza buchanani*, Pori, Finland, 31 October 2006 *(Harry Lehto)*

547 Hornemann's Redpoll / Groenlandse Witstuitbarmsijs *Carduelis hornemanni hornemanni*, Sumburgh Head, Mainland, Shetland, Scotland, 22 October 2006 *(Hugh Harrop)*

548 Lanceolated Warbler / Kleine Sprinkhaanzanger *Locustella lanceolata*, Fair Isle, Shetland, Scotland, 15 September 2006 *(Paul Baxter)*

24-25 October. If accepted, a female **'black-eared wheatear'** *O melanoleuca/hispanica* at Albarnaz lighthouse, Flores, on 9 October will be the first for the Azores. Until mid-November, the number of **Desert Wheatears** *O deserti* turning up in western Europe was lower than in recent years with, in Britain, first-winter females in Aberdeenshire on 21 October, on St Agnes, Scilly, on 27 October and at Cooden Beach, East Sussex, from 31 October to at least 4 November and a first-winter male on St Mary's from 1 November onwards. The first for western Turkey was found on Black Sea dunes near Karasu, Sakarya, 150 km from Istanbul, on 14 November. If accepted, a **Red-tailed Wheatear** *O xanthoprymna* near Konia in Paphos on 1 November will be the third for Cyprus. On 2 October, a **White's Thrush** *Zoothera aurea* was found on North Ronaldsay, Orkney, Scotland. In France, one was trapped on Hoëdic on 27 October. The second **Hermit Thrush** *Catharus guttatus* for Ireland was a first-winter on Cape Clear Island, Cork, on 19-20 October. On Ouessant, Finistère, two **Swainson's Thrushes** *C ustulatus* were found on 28 September, with one still present in early October. A first-winter female **Black-throated Thrush** *Turdus atrogularis* was on Foula on 7-9 October. In Scilly, **American Robins** *T migratorius* were seen on Tresco on 10-28 October and on Bryher on 23 October.

WARBLERS TO FLYCATCHERS **Pallas's Grasshopper Warblers** *Locustella certhiola* were trapped at Smokkevatnet, Time, Rogaland, on 24 September and on Whalsay on 2 October. Also in Shetland, **Lanceolated Warblers** *L lanceolata* were found on Fair Isle on 15 September (two) and on Foula on 5-9 October. The only WP record of the eastern subspecies of **Gray's Grasshopper Warbler** *L fasciolata amnicola* (split as **Sakhalin Grasshopper Warbler** *L amnicola* by Koblik et al in *Checklist of the birds of Russian Federation* 2006) on Ouessant on 17 September 1933 has been shown to be erroneous. It was listed as a juvenile *amnicola* in *Birds of the Western Palearctic* 6 (1992), but was rejected several years ago by the French rarities committee as it proved to be an adult nominate Gray's Grasshopper *L f fasciolata* in fresh plumage (which is impossible in September). A further examination of the specimen revealed that it concerns a missing specimen of the Natural History Museum in Tring, England, and that it belonged to the Owston collection from northeastern China in June 1908 (Br Birds 99: 506-516, 2006). As such, it constitutes another example of fraud by Richard Meinertzhagen involving stolen specimens which were incorporated in his own collection with false dates and locations (see, eg, Bull Br Orn Club 123A: 66-94, 2003, www.the-scientist.com/news/20051017/01). As *amnicola* is restricted to islands bordering the Sea of Okhotsk, the record was cited frequently as support for the occurrence of extreme vagrants from eastern Asia (eg, Dutch Birding 12: 9-11, 1990, 25: 221-234, 2003, 26: 53-56, 2004). There are two genuine records of juvenile nominate Gray's Grasshoppers collected as lighthouse victims, on Ouessant on 26 September 1913 and at Lodbjerg, Nordjylland, Denmark, on 25 September 1955. After two on 17 September, the third **Paddyfield Warbler** *Acrocephalus agricola* for the Netherlands this year was trapped at Meijendel, Zuid-Holland, on 14 October; there are now 18 records, all except one involving mistnetted birds. In Spain, one was trapped at Estanys de Palau, Aiguamolls de l'Empordà, on 9 October. In Cork, an **Eastern Olivaceous Warbler** *A pallidus* stayed on Cape Clear Island from 24 September to 1 October. The first **Cape Verde Warbler** *A brevipennis* for the northern half of Santiago was singing near Tarrafal on 17 October. In North Yorkshire, a **Two-barred Warbler** *Phylloscopus plumbeitarsus* was present at Filey on 16-18 October (Birding World 19: 435-436, 2006). A (very) late **Greenish Warbler** *P trochiloides* was photographed, sound-recorded and trapped at Zeebrugge, West-Vlaanderen, Belgium, on 6-7 November. A **Pallas's Leaf Warbler** *P proregulus* on Gambell, Alaska, on 25-26 September was the first for North America. By mid-October, more than 10 **Yellow-browed Warblers** *P inornatus* had arrived in Iceland. The third for North America was found on Attu, Alaska, on 21 September. The fourth **Radde's Warbler** *P schwarzi* for Italy was trapped on Ventotene in mid-October. In the Netherlands, three were found in Noord-Holland on 13 (trapped), 19 (trapped) and 29 October. On 12 October, a first-winter **Iberian Chiffchaff** *P ibericus* was photographed at Oukaimeden, High Atlas, Morocco, and several other individuals were present at this high-elevation ski resort as well. The first **Taiga Flycatcher** *Ficedula albicilla* for mainland North America was photographed at Putah Creek, Solano and Yolo, California, on 25 October. The first **European Pied Flycatcher** *F hypoleuca* for the Azores was found at Sete Cidades, São Miguel, on 27 October.

SHRIKES TO FINCHES The first **Red-backed Shrikes** *Lanius collurio* for the Azores were up to two juveniles on Corvo on 22-26 October. In the Netherlands, an adult male **Daurian Shrike** *L isabellinus* occurred at Den Hoorn, Texel, Noord-Holland, on 25-26 September. The first for Switzerland stayed at Laconnex, Genève, on 6-8 October. The second for Ireland was at Old Head of Kinsale, Cork, on 17-20 October. Another first-year was found at Cresswell Pond, Northumberland, on 28 October. The fifth for Spain was a first-year at Sanxenxo, Pontevedra, from 30 October to 1 November. In Scotland, a first-year **Turkestan Shrike** *L phoenicuroides* was present on Lewis, Outer Hebrides, on 22-29 September. The second **Long-tailed Shrike** *L schach* for Kuwait stayed at Sulaibikhat bay reserve from the last week of October to at least mid-November. An adult **Masked Shrike** *L nubicus* on Utsira on 2-15 October was the first for Norway. The second for Britain was a juvenile on St Mary's on 1 November. In England, **Red-eyed Vireos** *Vireo olivaceus* occurred at Nanquidno Valley, Cornwall, on 2-4 October, on Bryher on 17 October and on St Mary's on 19 October. In Ireland, one was at The Mullet, Mayo, on 5-7 October and another at Kilbaha,

549 Baltimore Oriole *Icterus galbula*, Cape Clear Island, Cork, Ireland, 17 October 2006 *(Paul & Andrea Kelly/irishbirdimages.com)* **550** Blackpoll Warbler / Zwartkopzanger *Dendroica striata*, first-year, Mandø, Denmark, 16 October 2006 *(Martin Søgaard Nielsen)* **551** Hermit Thrush / Heremietlijster *Catharus guttatus*, Cape Clear Island, Cork, Ireland, 19 October 2006 *(Paul & Andrea Kelly/irishbirdimages.com)* **552** Yellow-breasted Bunting / Wilgengors *Emberiza aureola*, Whalsay, Shetland, Scotland, 11 September 2006 *(Hugh Harrop)*

Clare, on 7-14 October. Three single **European Goldfinches** *Carduelis carduelis* at three sites during 12-16 October were the second to fourth for Iceland (the first was on 17 October 2005); a fifth was found on 1 November. A **Greenland Redpoll** *C flammea rostrata* on Corvo on 21 October was the second redpoll for the Azores and Macaronesia. In Norway, one was found at Mønstremyr, Vest-Agder, on 2 November. Two **Hornemann's Redpolls** *C hornemanni hornemanni* were reported from Røst, Nordland, Norway, on 2 October. Four singles occurred in Shetland on Foula on 4-13 October, at Sumburgh on 19-24 October and on Fair Isle on 25 and 30 October. Recent sound-recordings of **British Crossbill** *Loxia curvirostra* type E in Estonia were the first outside Britain and the Netherlands; perhaps less surprising were the first sound-recordings of **Scarce Crossbills** *L c* type F outside the Benelux in Estonia and Russia (cf *The Sound Approach to birding* 2006). In coastal south-western Norway, an influx of **Pine Grosbeak** *Pinicola enucleator* became apparent in early November, with a flock of 109 on 11 November.

AMERICAN WARBLERS TO AMERICAN BLACKBIRDS In Ireland, a **Blackpoll Warbler** *Dendroica striata* turned up on Dursey Island, Cork, on 10 October. The first for Denmark was an immature female on Mandø in the Waddensea south of Esbjerg, Vestjylland, on 15-16 October and the first for Norway was at Reve, Klepp, Rogaland, from 2 to at least 11 November. A first-winter male **Common Yellowthroat** *Geothlypis trichas* was found dead as a window victim at Penryn, Cornwall, on 23 October. The first **Canada Warbler** *Wilsonia canadensis* for Ireland was a first-winter female at Kilbaha near Loop Head on 8-13 October (Birding World 19: 429-434, 2006); the only previous WP record was a male in Iceland on 29 September 1973. A male **American Redstart** *Setophaga ruticilla* flying in front of a driving car at La Janda, Cadiz, and

553 Canada Warbler / Canadese Zanger *Wilsonia canadensis*, Kilbaha, Clare, Ireland, October 2006 *(John Carter)*

554 Blackpoll Warbler / Zwartkopzanger *Dendroica striata*, first-year, Reve, Klepp, Rogaland, Norway, 2 November 2006 *(Christian Tiller)*

subsequently seen for a few minutes on 6 October was, if accepted, the first for Spain; the most recent WP records were in Ireland and Britain in October 1985. A **Lapland Longspur** *Calcarius lapponicus* photographed at Rumeli Feneri, Istanbul, on 2 November was the first for Turkey. In Norway, a female **Pine Bunting** *Emberiza leucocephalos* was seen at Reime, Hå, Rogaland, on 21 October. Yet another reintroduction in Britain and the first of a passerine species involved more than 70 **Cirl Buntings** *E cirlus* in southern Cornwall. These birds were taken from the nest as tiny chicks this summer in southern Devon and were hand-reared before being released. A **Grey-necked Bunting** *E buchanani* seen by a handful of birders at Pori on 30 October was first identified as Ortolan Bunting *E hortulana* before it was re-identified after sunset; the next day, it was still present and seen by 100s of twitchers as the first for Finland and the second for western Europe (the first was trapped at Castricum, Noord-Holland, on 16 October 2004). **Yellow-breasted Buntings** *E aureola* were on Whalsay on 11 September and on Fair Isle on 15-16 September. The fifth **Pallas's Reed Bunting** *E pallasi* for Alaska and North America was on Gambell on 24-26 September. In Britain, male **Black-headed Buntings** *E melanocephala* were seen at Holme, Norfolk, on 13 September, at Toab, Shetland, on 16-19 September and on Fetlar, Shetland, on 22-24 September; the 12th for the Netherlands (and first in autumn) occurred at Brouwersdam, Zeeland, on 16-18 October. A **Baltimore Oriole** *Icterus galbula* on Cape Clear Island on 12-19 October was the second for Ireland.

NEARCTIC VAGRANTS IN THE AZORES Like last year, an amazing number of Nearctic vagrants was seen in the Azores (cf Dutch Birding 27: 424-425, 2005); by late October, they were found by 12 birders on Corvo and 26 on Flores. For **Pale-bellied Brent Geese**, see above. **Lesser Scaups** *A affinis* were found at Pedro Miguel, Faial, on 1 November (first-winter female), at Sete Cidades, São Miguel, on 3 November (six), at Lagoa Azul, on 5 November and at Lagoa dos Patos, on 4 November (female). On Flores, seven **Ring-necked Ducks** *A collaris*, a first-year female **American Wigeon** *A americana*, two juvenile **Blue-winged Teals** *A discors*, two **American Black Ducks** with five hybrids, and 16 juvenile (presumed) **Green-winged Teals** *A carolinensis* were found at Lagoa Seca on 7-8 November. A few American Wigeons and Ring-necked Ducks were also seen on São Miguel, Terceira, São Jorge and/or Pico. During October, one **American Black Duck** was also present on São Miguel and up to 15, including hybrids, were on Corvo. The long-staying male **Wood Duck** *Aix sponsa* was still swimming at a park at Furnas, São Miguel, on 5 November. A juvenile **Double-crested Cormorant** *P auritus* was present at Mosteiros, São Miguel, from 25 October into November. A **Green Heron** was observed at Fajã Grande between 22 October and 2 November. An adult and a juvenile **Snowy Egret** *E thula* were seen at Fajã Grande from 3 November. **American Great Egrets** *C a egretta* were present on São Miguel on 30 September and 1 October, on Terceira on 19 October, and at Fajã Grande, Flores, from 31 October to at least 8 November. On Flores, the long-staying **Great Blue Heron** *Ardea herodias* from 22 September 2005 was still present at Lagoa Branca on 7 November. From 17 October into November, at least four other individuals turned up on Flores. A juvenile **Pied-billed Grebe** *Podilymbus podiceps* was discovered on Corvo on 29 October. An **American Coot** *Fulica americana* remained at Lagoa das Furnas, São Miguel, until at least 5 November, while another was found at Lagoa do Caiado, Pico, on 2 November. A juvenile **Killdeer** was on Corvo on at least 27-29 October. On Terceira, the following Nearctic waders were seen at Cabo da Praia from 15 September to at least 4 November: up to two **Semipalmated Plovers**, five **American Golden Plovers**, four **Semipalmated Sandpipers**, a **Western Sandpiper**, a **Least Sandpiper**, c 18 **White-rumped Sandpipers** *C fuscicollis*, a **Baird's Sandpiper** *C bairdii*, three **Pectoral Sandpipers**, a **Short-billed Dowitcher** *L griseus*, two to four **Long-billed Dowitchers** (first records), five **Spotted Sandpipers** *Actitis macularius*, a **Greater Yellowlegs** *T melanoleuca* and four **Lesser Yellowlegs** *T flavipes*. Some of these species were found on other islands as well. **Least Sandpipers** were also present at Ponta Delgada on 13 September and at Lajes do Pico, Pico, on 19-20 September. The ninth **Wilson's Snipe** *Gallinago delicata* for the Azores was reported in a flock of 75 Common Snipes *G gallinago* at Caldeirão, Corvo, on 6 October. **Short-billed Dowitchers** were also seen at Lajes do Pico on 22 September and 2 November. On 5 October, a **Hudsonian Whimbrel** *Numenius hudsonicus* turned up north of Praia da Vitória, Terceira. An **Upland Sandpiper** was seen at Ponta Delgada from 29 September to 8 October. The third and fourth **Willet** *Tringa semipalmata* were reported at Fajã dos Cubres, São Jorge, on 22 September and at Ponta Delgada on 4 November. Two **Laughing Gulls** were present at Cabo da Praia on 21 September. A first-winter **American Herring Gull** *L smithsonianus* was identified at Ponta Delgada on 3-5 November. On Flores, the fourth **Black-billed Cuckoo** *Coccyzus erythropthalmus* occurred at Fajã Grande on 28-29 October and a **Yellow-billed Cuckoo** *C americanus* was discovered also at Fajã Grande on 26 October, with another at Ponta Delgada on 1 November. **Chimney Swifts** *Chaetura pelagica* were seen on Corvo on 29 and 31 October and 2 November. On Flores, one flew about at Santa Cruz on 31 October, two at Ladeira da Picada on 1 November and one at Fajã Grande on 1 November. Two first-year **American Barn Swallows** *Hirundo rustica erythrogaster* on Corvo on 20-29 October constituted the third for the WP (cf Dutch Birding 28: 44, plate 47, 2006). **Red-eyed Vireos** on Corvo on 23-25 and 28 October were (only) the second and third. A first-winter female **Black-throated Blue Warbler** *D caerulescens* on Corvo on 24 and 28-29 October and a male on 29 October were also the second and third. Three single first-winter **Myrtle Warblers** *D coronata* on Corvo on 21, 25 and

26 October were the second to fourth. A **Blackpoll Warbler** at Aldeia da Cuada, Flores, on 24 October was (only) the first. The second **Common Yellowthroat** *Geothlypis trichas* was a male at Fajázinha, Flores, on 20-21 October. The first **Summer Tanager** *Piranga rubra* (and the second for the WP) was found on Corvo on 26 October and seen again on 28 October; the first for the WP was on Bardsey, Wales, in September 1957. An adult **Lapland Longspur** on Corvo on 21 October could be either of Nearctic or Palearctic origin. Also on Corvo, a female **Rose-breasted Grosbeak** *Pheucticus ludovicianus* was seen on 23 October and a male and a female on 24 October. First-year **Indigo Buntings** *Passerina cyanea* were found on 24, 27 and 29 October and 1 November on Corvo, and other first-years were seen at Aldeia da Cuada on 24-25 October and at Fajã Grande on 28 October.

For a number of reports, Birding World, Birdwatch, British Birds, Ornithos, www.azores.seawatching.net, www.birdguides.com, www.netfugl.dk and www.rarebirdalert.co.uk were consulted. We wish to thank Peter Alfrey (Azores), Chris Batty, Soner Bekir (Turkey), Morten Bentzon Hansen, Max Berlijn, Keith Betton, Richard Bonser, Rob Brouwer (Cape Verde Islands), David Camilleri, Rommert Cazemier, Sean Cole, Andrea Corso (Italy), Bram Couperus, Pierre-André Crochet, Kris De Rouck, Jochen Dierschke, J Ignacio Dies, Hugues Dufourny, Enno Ebels, Lee Evans, Pekka Fagel, Frode Falkenberg, Dick Forsman, Raymond Galea (Malta), Steve Gantlett, Arthur Geilvoet, Jeff Gordon, Barak Granit, George Gregory (Kuwait), Marcello Grussu (Sardinia), Ricard Gutiérrez (Spain), Carlos Gutiérrez Expósito, Hugh Harrop, Paul Harvey, Cornelis Hazevoet, Erik Hirschfeld, Niklas Holmström, Erling Jirle (Sweden), Adrian Jordi, Peter Kennerley, Howard King, Szabolcs Kókay, Yann Kolbeinsson, Tomek Kulakowski, Henning Kunze, Joep de Leeuw, Antero Lindholm (crossbills), André van Loon, Erik Maassen, Guy McCaskie, Peter Meijer, Richard Millington, Dominic Mitchell, Nial Moores, Richard Moores, Killian Mullarney, Daniele Occhiato, Gert Ottens, Benoit Paepegaey, Menotti Passarella, Andy Paterson, Yoav Perlman (IRDC), Ron Pittaway, René Pop, Mike Pope, Carla Rapetti, Colin Richardson, Matthias Ritschard, Magnus Robb, Michel Rogg, René van Rossum, Luciano Ruggieri, Michael Sammut, Nir Sapir, Dare Šere, Jean-Philippe Siblet, Sébastien Siblet, Russell Slack (BirdGuides), Roy Slaterus, Brian Small, Michael Smith, Darryl Spittle, John Sterling, Roland van der Vliet, Bernard Volet, Damian Wiehle, Edwin Winkel (Cape Verde Islands), Rik Winters, Emin Yogurtcuoglu and Steve Young for their help in compiling this review.

Arnoud B van den Berg, Duinlustparkweg 98, 2082 EG Santpoort-Zuid, Netherlands (arnoud.vandenberg@planet.nl)
Marcel Haas, Hora Siccamasingel 167, 9721 HD Groningen, Netherlands (m.haas@wpbirds.com)

Recente meldingen

Dit overzicht van recente meldingen van zeldzame en interessante vogels in Nederland en België beslaat voornamelijk de periode **september-oktober 2006**. De vermelde gevallen zijn merendeels niet geverifieerd en het overzicht is niet volledig. Alle vogelaars die de moeite namen om hun waarnemingen aan ons door te geven worden hartelijk bedankt. Waarnemers van soorten in Nederland die worden beoordeeld door de Commissie Dwaalgasten Nederlandse Avifauna wordt verzocht hun waarnemingen zo spoedig mogelijk toe te zenden aan: CDNA, Postbus 45, 2080 AA Santpoort-Zuid, Nederland, e-mail cdna@dutchbirding.nl. Hiertoe gelieve men gebruik te maken van CDNA-waarnemingsformulieren die eveneens verkrijgbaar zijn bij bovenstaand adres, of via de website van de DBA op www.dutchbirding.nl.

Nederland

GANZEN TOT VALKEN Na de eerste melding van twee **Dwergganzen** *Anser erythropus* op 29 september aan de oostkant van de Lauwersmeer, Groningen, werd op 14 oktober al een groep van 37 gemeld bij de Anjumer Kolken, Friesland, en op 29 oktober een groep van 17 in het Oude Land van Strijen, Zuid-Holland. Daarnaast werden nog c 10 losse exemplaren gemeld. Een **Groenlandse Kolgans** *A albifrons flavirostris* werd van 12 tot 14 oktober gezien bij Nieuw-Scheemda, Groningen. Na de eerste melding van een **Roodhalsgans** *Branta ruficollis* op 23 september bij Aldegea (Oudega) in Gaasterland, Friesland, werden er vanaf 6 oktober nog eens 12 gezien verspreid over het land. Vanaf begin oktober werden zowel zes **Witbuikrotganzen** *B hrota* als zes **Zwarte Rotganzen** *B nigricans* gemeld. De beste plekken voor **Casarca's** *Tadorna ferruginea* waren de Vreugderijkerwaard, Overijssel, met 121 op 15 september en het Ketelmeer, Overijssel, met 60 op 30 september. Grote aantallen **Krooneenden** *Netta rufina* zwommen op 28 september op het Veluwemeer bij Nunspeet, Gelderland (84), op 14 oktober op de Gouwzee, Noord-Holland (120), en op 26 oktober niet minder dan 375 op een plas in het Harderbroek, Flevoland. Het mannetje **Grote Tafeleend** *Aythya valisineria* van het Hoefijzermeer bij Castricum, Noord-Holland, was daar weer aanwezig vanaf 30 september. Vanaf 14 september werd een 10-tal **Witoogeenden** *A nyroca* gemeld. Op 1 september zwom het vrouwtje **Ringsnaveleend** *A collaris* nog bij Budel-Dorplein, Noord-Brabant. Adulte mannetjes **Buffelkopeend** *Bucephala albeola* werden gezien op 17 september op het IJmeer bij Muiden, Noord-Holland, van 24 september tot 5 oktober op de Gouwzee, Noord-Holland, vanaf 13 oktober weer bij Muiden en op 23 oktober ook op het Naardermeer, Noord-Holland. **IJsduikers** *Gavia immer* vlogen op 4 oktober langs Vlieland, Friesland, op 7 oktober langs Rottumeroog, Groningen, op 25 oktober langs IJmuiden, Noord-Holland, en op 31 oktober langs Westkapelle, Zeeland.

555 Roodhalsganzen / Red-breasted Geese *Branta ruficollis*, met Rotganzen / Dark-bellied Brent Geese *B bernicla*, Terschelling, Friesland, 8 oktober 2006 *(Arie Ouwerkerk)*

Grauwe Pijlstormvogels *Puffinus griseus* waren schaars met als uitschieters bijvoorbeeld 36 op 7 oktober langs Texel, Noord-Holland, en 29 op 24 oktober langs Westkapelle. Tussen 26 september en 29 oktober werden slechts 30 **Noordse Pijlstormvogels** *P puffinus* doorgegeven. **Vale Pijlstormvogels** *P mauretanicus* werden vaker gemeld dan normaal met waarnemingen op 4 september langs Westkapelle, op 16 september langs Camperduin, Noord-Holland, op 26 september langs Scheveningen, Zuid-Holland, op 4 oktober langs Terschelling, Friesland, en op 7 oktober langs Texel, de Maasvlakte, Zuid-Holland, en Vlieland. **Stormvogeltjes** *Hydrobates pelagicus* werden opgemerkt op 4 oktober langs Terschelling en op 31 oktober langs Westkapelle. Slechts enkele **Vale Stormvogeltjes** *Oceanodroma leucorhoa* werden gemeld, met een uitzonderlijke waarneming op 18 oktober over de Loozerheide, Noord-Brabant. Dit exemplaar werd in vlucht gefotografeerd en pas achteraf door de verbaasde waarnemers gedetermineerd. **Kuifaalscholvers** *Phalacrocorax aristotelis* werden gezien tot 8 oktober bij IJmuiden, op 1 september bij Den Helder, Noord-Holland, en op 13 en 20 oktober bij de Oosterscheldekering, Zeeland. Naar wordt aangenomen dezelfde **Groene Reiger** *Butorides virescens* als die in april-juni verbleef bij De Nieuwe Meer bij Amsterdam, Noord-Holland, werd van 14 tot 30 september (volgens lokale informatie al vanaf juli) met enige regelmaat gezien bij de Noorder IJ-plas bij Amsterdam; de vogel was nu in onvolwassen zomerkleed. **Koereigers** *Bubulcus ibis* verschenen op 12 september bij Groot-Ammers, Zuid-Holland, op 30 september in De Putten en in de Hilversumsche Bovenmeent, Noord-Holland, op 9 oktober in de Ooypolder, Gelderland, en vanaf 16 oktober bij Kruispolderhaven, Zeeland. Een aardige verzameling van 62 **Grote Zilverreigers** *Casmerodius albus* verbleef op 17 oktober in Polder Maltha in de Brabantsche Biesbosch, Noord-Brabant. Net als met de Grote Zilverreiger gaat het ook met de **Kleine Zilverreiger** *Egretta garzetta* als broedvogel voorspoedig. Dit jaar omvatte de totale populatie 125 paren, met alleen al 55 paren in de kolonie van het Quackjeswater, Zuid-Holland, 25 in Zeeuws-Vlaanderen, Zeeland, en 25 in de Oostvaardersplassen, Flevoland. Op 12 augustus werd de afwijkende donkere **Blauwe Reiger** *Ardea cinerea* met kromme snavel die tot begin augustus in Capelle aan den IJssel, Zuid-Holland, verbleef gefotografeerd in Alblasserdam, Zuid-Holland (cf Dutch Birding 28: 299-300, 2006). In september werden langs De Gorzen bij Ridderkerk, Zuid-Holland, c 250 langsvliegende **Purperreigers** *Ardea purpurea* geteld. Er werden vele 10-tallen **Zwarte Ooievaars** *Ciconia nigra* gezien met als grootste aantallen zeven op 1 september over de Strabrechtse Heide, Noord-Brabant, vijf op 16 september over Benthuizen, Zuid-Holland, en drie tot vier in de tweede helft van september in De Banen bij Nederweert, Limburg. **Zwarte Ibissen** *Plegadis falcinellus* verschenen op 4 september in De Bijland, Gelderland, op

556 Blonde Ruiter / Buff-breasted Sandpiper *Tryngites subruficollis*, juveniel, De Stolpen, Noord-Holland, 28 september 2006 *(Roland Jansen)* **557** Blonde Ruiter / Buff-breasted Sandpiper *Tryngites subruficollis*, juveniel, De Stolpen, Noord-Holland, 27 september 2006 *(Leo J R Boon/Cursorius)* **558** Blonde Ruiter / Buff-breasted Sandpiper *Tryngites subruficollis*, juveniel, Slikken van Flakkee, Zuid-Holland, 13 september 2006 *(Pim A Wolf)*

559 Blonde Ruiter / Buff-breasted Sandpiper *Tryngites subruficollis*, juveniel, De Stolpen, Noord-Holland, 27 september 2006 *(Garry Bakker)*

560 Blonde Ruiters / Buff-breasted Sandpipers *Tryngites subruficollis*, juveniel, Amaliaweg, Texel, Noord-Holland, 25 september 2006 *(René Pop)*

561 Hop / Eurasian Hoopoe *Upupa epops*, Castricum, Noord-Holland, 9 oktober 2006
(Ton Nagtegaal)

562 Nachtzwaluw / European Nightjar *Caprimulgus europaeus*, Kwintsheul, Zuid-Holland, 16 september 2006
(Leo J R Boon/Cursorius)

563 Roodkopklauwier / Woodchat Shrike *Lanius senator*, eerstejaars, De Cocksdorp, Texel, Noord-Holland, 2 september 2006 *(Harvey van Diek)* **564** Zwartkopgors / Black-headed Bunting *Emberiza melanocephala*, mannetje, Brouwersdam, Zeeland, 17 oktober 2006 *(Leo J R Boon/Cursorius)* **565** Daurische Klauwier / Daurian Shrike *Lanius isabellinus*, adult mannetje, Den Hoorn, Texel, Noord-Holland, 26 september 2006 *(Rein Hofman)*

566 Grote Tafeleend / Canvasback *Aythya valisineria*, adult mannetje, Hoefijzermeer, Castricum, Noord-Holland, 30 september 2006 *(Martijn Verdoes)* **567** Klein Waterhoen / Little Crake *Porzana parva*, Verdronken Land van Saeftinghe, Zeeland, 8 september 2006 *(Alex Wieland)* **568** Groene Reiger / Green Heron *Butorides virescens*, tweede-kalenderjaar, Zijkanaal H-Weg, Amsterdam, Noord-Holland, 19 september 2006 *(Effendi Haidar)* **569** Groene Reiger / Green Heron *Butorides virescens*, tweede-kalenderjaar, Zijkanaal H-Weg, Amsterdam, Noord-Holland, 30 september 2006 *(Enno B Ebels)*

6 september over Veenendaal, Utrecht, op 1 oktober bij Camperduin, op 5 oktober twee (overvliegend) bij Lexmond, Zuid-Holland, op 8 oktober over Texel, en op 9 oktober langs Katwijk aan Zee, Zuid-Holland. Uiteraard was ook de vogel van het Balgzand en omstreken, Noord-Holland, de gehele periode aanwezig. Er werd een achttal **Zwarte Wouwen** *Milvus migrans* gemeld waaronder één van 4 tot 6 september in het Markiezaat, Noord-Brabant, soms in gezelschap van twee **Rode Wouwen** *M milvus*. Tot half oktober trokken c 10 **Zeearenden** *Haliaeetus albicilla* het land in. Het broedpaar met hun jonge vogel bleef aanwezig in de Oostvaardersplassen en vanaf 16 oktober was er één aanwezig in de wijde omgeving van de Korendijkse Slikken, Zuid-Holland. De **Slangenarend** *Circaetus gallicus* van Meijendel, Zuid-Holland, bleef tot 9 september. Een mogelijke **Steppebuizerd** *Buteo buteo vulpinus* werd op 27 oktober gefotografeerd bij Lettele, Overijssel, en een **Arendbuizerd** *B rufinus* passeerde op 29 september de telpost Kamperhoek, Flevoland. Er waren meldingen van **Dwergarenden** *Aquila pennata* op 8 september bij Nieuwegein, Utrecht, en op 5 oktober bij Hoogvliet, Zuid-Holland. Spectaculair was de waarneming van twee subadulte **Steenarenden** *A chrysaetos* op 16 september langs de telpost IJmeerdijk bij Almere, Flevoland. Alleen al in september werden c 50 **Roodpootvalken** *Falco vespertinus* gemeld. In oktober volgden er nog drie met de laatste op 16 oktober bij Heemstede, Noord-Holland. Een ontsnapte grote valk met een blauwe ring aan de linkerpoot werd op 15 oktober gefotografeerd op de Westplaat, Zuid-Holland, en vloog twee weken later bij de Stuifdijk op de Maasvlakte; hij leek sterk op een Giervalk *F rusticolus* maar vertoonde hybride invloed van waarschijnlijk Sakervalk *F cherrug*.

570 Raddes Boszanger / Radde's Warbler *Phylloscopus schwarzi*, Kennemerduinen, Bloemendaal, Noord-Holland, 13 oktober 2006 *(Erik Maassen/Vrs van Lennep)* **571** Veldrietzanger / Paddyfield Warbler *Acrocephalus agricola*, Meijendel, Zuid-Holland, 14 oktober 2006 *(Vincent van der Spek/Vrs Meijendel)* **572** Siberische Boompieper / Olive-backed Pipit *Anthus hodgsoni*, Oosterreeweg, Schiermonnikoog, Friesland, 15 oktober 2006 *(Martijn Bot)* **573** Kortteenleeuwerik / Greater Short-toed Lark *Calandrella brachydactyla*, Oosterend, Terschelling, Friesland, 25 oktober 2006 *(Arie Ouwerkerk)*

RALLEN TOT ALKEN Op 8 september werd een eerstejaars **Klein Waterhoen** *Porzana parva* gevangen en geringd in het Verdronken Land van Saeftinghe, Zeeland. **Kraanvogels** *Grus grus* trokken door vanaf 12 oktober en vooral op 16 oktober werden grote groepen waargenomen, zoals ten minste 113 over Epen, Limburg, en meer dan 1000 over Maastricht, Limburg. Een **Steltkluut** *Himantopus himantopus* werd op 7 oktober gemeld op het Landje van Naber bij Hoorn, Noord-Holland. De **Steppevorkstaartplevier** *Glareola nordmanni* van de omgeving van Ransdorp werd weer gezien van 13 tot 18 september. Van 14 tot 20 oktober verbleef een exemplaar bij Eemnes, Utrecht. Ongedetermineerde **vorkstaartplevieren** vlogen op 17 oktober over natuurontwikkelingsgebied Willeskop, Utrecht, en bij telpost Brobbelbies bij Uden, Noord-Brabant. Er werden tot 13 oktober nog c 70 **Morinelplevieren** *Charadrius morinellus* gemeld. Vermeldenswaard zijn twee exemplaren aan de grond op 2 september op De Hoge Veluwe, Gelderland, en drie op 7 september bij Diependal, Drenthe, en 24 over de telpost in de Groote Peel op 8 september. Er waren meldingen van een **Aziatische Goudplevier** *Pluvialis fulva* op 16 september bij Nieuwstad, Groningen, en van een **'kleine goudplevier'** *P dominica/fulva* op 16 oktober over polder Kwistenburg op Zuid-Beveland, Zeeland. Erg leuk is de waarneming van een **Steppekievit** *Vanellus gregarius* op 15 september bij Budel-Dorplein. De vogel vloog later die dag richting België. Een **Bonapartes Strandloper** *Calidris fuscicollis* werd op 13 oktober kortstondig gezien aan het Amstelmeer, Noord-Holland. Pas in deze periode nam het aantal waarnemingen van **Gestreepte Strandlopers** *C melanotos* sterk toe tot ten minste 25, waaronder zeker vier op 17 september in de Lauwersmeer en drie van 22 tot 25 september in de Scherpenissepolder, Zeeland. **Breedbekstrandlopers** *Limicola falcinellus* werden waargeno-

574 Vale Gierzwaluw / Pallid Swift *Apus pallidus*, Vlieland, Friesland, 20 oktober 2006 *(Bas van den Boogaard)* **575** Steppevorkstaartplevier / Black-winged Pratincole *Pratincola nordmanni*, Eemnes, Utrecht, 17 oktober 2006 *(Harm Niesen)* **576** Roodpootvalk / Red-footed Falcon *Falco vespertinus*, juveniel, Westkapelle, Zeeland, 30 september 2006 (*Corstiaan Beeke)*

577 Kleine Spotvogel / Booted Warbler *Acrocephalus caligatus*, Maasvlakte, Zuid-Holland, 9 oktober 2006 *(Roland Jansen)*

578 Kleine Spotvogel / Booted Warbler *Acrocephalus caligatus*, Maasvlakte, Zuid-Holland, 9 oktober 2006 *(Jan den Hertog)*

579 Cetti's Zanger / Cetti's Warbler *Cettia cetti*, Dordtse Biesbosch, Zuid-Holland, 10 oktober 2006
(Hans Gebuis)

580 Raddes Boszanger / Radde's Warbler *Phylloscopus schwarzi*, Castricum, Noord-Holland, 19 oktober 2006
(Cock Reijnders)

men op 1 september bij Zwarte Haan, Friesland, op 9 september (drie!) bij Paesens, Friesland, op 14 september bij Holwerd, Friesland, van 15 tot 20 september in Polder IJdoorn, Noord-Holland, en op 30 september in Polder Breebaart, Groningen. Er was sprake van een influx van **Blonde Ruiters** *Tryngites subruficollis* met exemplaren op 9 september op Vlieland, op 13 september op de Slikken van Flakkee, Zuid-Holland, en op 21 september in Polder de Bant, Friesland; op 24 september werden er zelfs twee bij elkaar gevonden op Texel en één bij De Stolpen, Noord-Holland, die respectievelijk tot 28 en 29 september bleven. In alle gevallen betrof het juveniele vogels. Een **Poelsnip** *Gallinago media* was op 12 en 13 september aanwezig bij Zwolle, Overijssel, in hetzelfde gebied (Hessenpoort) als vorig jaar op 13 september. Op 16 september werd een exemplaar ontdekt op Texel. Een **Kleine Geelpootruiter** *Tringa flavipes* werd kort waargenomen op 9 september bij Holwerd en pas gedetermineerd aan de hand van een foto. **Poelruiters** *T stagnatilis* verbleven van 8 tot 13 september in de Jan Durkspolder bij Oudega, Friesland, van 10 tot 28 september (maximaal twee) in de Lauwersmeer, op 22 september bij De Schelphoek, Zeeland, op 24 september bij Stavenisse, Zeeland, op 6 oktober op Terschelling en op 23 oktober bij Retranchement, Zeeland. Na een vroege **Rosse Franjepoot** *Phalaropus fulicarius* op 10 september bij IJmuiden kwamen er 15 in oktober waarvan zeven tussen 4 en 8 oktober en zes op 29 oktober. Een pleisteraar verbleef van 20 tot 27 oktober op Terschelling. Tot 9 oktober werden nog c 30 **Kleinste Jagers** *Stercorarius longicaudus* waargenomen, waaronder één van 3 tot 8 september bij Goudswaard, Zuid-Holland, en vijf op 3 oktober langs Westkapelle. Vanaf 6 september werden bijna 20 **Vorkstaartmeeuwen** *Larus sabini* gezien waarvan het tweetal dat op 4 oktober over telpost Kwintelooijen bij Veenendaal vloog extra het vermelden waard is. Een adulte **Dunbekmeeuw** *L genei* werd op 14 september gemeld en op de plaat gezet bij Veessen, Overijssel. Op 16 september werden er vier gemeld langs de Nolledijk bij Vlissingen, Zeeland. Een mogelijke (subadulte) **Amerikaanse Zilvermeeuw** *L smithsonianus* werd op 27 oktober gefotografeerd bij Westkapelle en verdween kort daarna in zuidelijke richting. De **Grote Burgemeester** *L hyperboreus* van Katwijk aan Zee hield de gehele periode voet bij stuk. De adulte van Den Helder was weer terug vanaf 6 september en een andere adulte vloog op 29 oktober zuidwaarts langs de Langevelderslag, Noord-Holland. De **Ringsnavelmeeuw** *L delawarensis* van Tiel, Gelderland, werd regelmatig gezien. **Lachsterns** *Gelochelidon nilotica* werden waargenomen op 5 september in de Oostvaardersplassen, op 6 september bij Rottumeroog, Groningen, op 17 en 23 september bij Lauwersoog, Groningen, en op 23 september roepend in een groep Grote Sterns *Sterna sandvicensis* bij IJmuiden. De hoogste aantallen **Reuzensterns** *Hydroprogne caspia* werden geteld op 9 september in de Makkumerzuidwaard, Friesland (28), en op 14 september in de Workumerwaard, Friesland (33). Late exemplaren werden op 8 oktober opgemerkt bij IJmuiden en de Oostvaardersdijk. Een **Witwangstern** *Chlidonias hybrida* trok op 27 september langs Scheveningen. Nog c 20 **Witvleugelsterns** *C leucopterus* werden waargenomen, voornamelijk in september. Het langst bleef een exemplaar bij Stellendam (tot 10 oktober) en op 14 oktober vloog er één over Texel. **Zwarte Zeekoeten** *Cepphus grylle* werden gezien op 7 oktober (twee) en 15 oktober (één) langs Vlieland, en op 9 oktober bij Texel. Al op 5 oktober vloog een **Kleine Alk** *Alle alle* langs Scheveningen. Tussen 24 oktober en 1 november werden er aan de kust c 100 gezien met op 25 oktober 16 langs Ameland, Friesland, en 32 langs Schiermonnikoog. **Papegaaiduikers** *Fratercula arctica* vlogen op 8 en 22 oktober langs Vlieland, op 13 oktober langs Terschelling, op 25 oktober langs Schiermonnikoog, Friesland, en op 29 oktober langs Texel.

NACHTZWALUWEN TOT GORZEN Een **Nachtzwaluw** *Caprimulgus europaeus* verbleef van 13 tot 17 september overdag in een tuin van een vogelaar te Kwintsheul, Zuid-Holland, en trok nogal wat bekijks. Eind oktober werden c 10 **Gierzwaluwen** *Apus apus* gezien. Op 20 oktober gingen er twee op Vlieland (eindelijk) vergezeld van één **Vale Gierzwaluw** *A pallidus*. Op 21 oktober bleek deze vermoedelijk eerste voor het land nog aanwezig en kon, tot groot genoegen van de overgekomen vogelaars, tot in de avondschemering met enige regelmaat worden gezien. Op 22 oktober werd hij helaas niet teruggevonden. Op 25 oktober waren er nog meldingen van deze soort over De Groote Peel, Noord-Brabant, en langs de Nolledijk bij Vlissingen. Redelijk laat waren de vier **Bijeneters** *Merops apiaster* op 4 september in het gebied Drentsch Friesche bij Westerveld, Drenthe. Op 6 september vlogen er twee over Vasse, Overijssel. **Hoppen** *Upupa epops* verbleven tot 1 september bij Linden, Noord-Brabant, van 31 augustus tot 5 september en op 19 september bij Bergen aan Zee, Noord-Holland, op 2 september in Roermond, Limburg, op 11 september in Arnhem, Gelderland, op 16 september in de AW-duinen, Noord-Holland, en van 5 tot 8 oktober bij Castricum. Een late **Draaihals** *Jynx torquilla* werd op 3 oktober geringd op Schiermonnikoog. Er waren **Kortteenleeuweriken** *Calandrella brachydactyla* op 2 oktober in de Eemshaven, Groningen, en van 15 tot maar liefst 25 oktober bij Oosterend op Terschelling. Op 23 september trok een **Roodstuitzwaluw** *Cecropis daurica* over telpost De Vulkaan bij Den Haag, Zuid-Holland. Het optreden van de **Grote Pieper** *Anthus richardi* verliep ongeveer als volgt: tussen 9 en 16 september drie, van 22 tot 30 september minstens 13 en in oktober zeker c 50 exemplaren. Er werden c 65 **Duinpiepers** *A campestris* doorgegeven, voornamelijk in september, met hekkensluiters op 1 oktober over het Leersumse Veld, Utrecht, en op 3 oktober langs De Hondsbossche Zeewering bij Camperduin. Een **Siberische Boompieper** *A hodgsoni* was op 15 en 16 oktober aanwezig in het dorp op Schiermonnikoog. De eerste **Roodkeelpieper** *A cervinus* van het najaar werd op 12 september waargenomen bij Veenendaal, spoedig gevolgd door enkele 10-tallen tot 16 oktober en een laat exemplaar op 29 oktober bij Camperduin. De eerste **Pestvogel** *Bombycilla garrulus* dit najaar vloog op

581 Zwartkopgors / Black-headed Bunting *Emberiza melanocephala*, mannetje, Brouwersdam, Zeeland, 17 oktober 2006 *(Roland Jansen)*

582 Bosgors / Rustic Bunting *Emberiza rustica*, Vlieland, Friesland, 7 oktober 2006 *(Bas van den Boogaard)*

8 oktober over de Eemshaven. Er waren nog twee ringvangsten van **Noordse Nachtegaal** *Luscinia luscinia*, op 17 september op Schiermonnikoog en op 21 september te Bloemendaal, Noord-Holland. Een **Aziatische Roodborsttapuit** *Saxicola maurus* werd op 8 oktober gemeld op Texel. De **Izabeltapuit** *Oenanthe isabellina* die vanaf 31 augustus in de Eemshaven verbleef werd na 4 september niet meer gezien. **Cetti's Zangers** *Cettia cetti* verbleven tot 16 september bij Zwijndrecht, Zuid-Holland, (één of twee) tot 27 oktober bij Stellendam, Zuid-Holland, op 17 september bij IJmuiden, op 26 september bij Breskens, Zeeland, van 8 tot 21 oktober nog steeds bij Werkendam, Zuid-Holland, van 10 tot 22 oktober bij telpost Dordtse Biesbosch, Zuid-Holland, en op 21 oktober nabij de Brouwersdam, Zuid-Holland. Op 15 oktober werd de derde dit jaar geringd voor het Verdronken Land van Saeftinghe. In september werden ruim 30 **Graszangers** *Cisticola juncidis* doorgegeven, voornamelijk uit Zuid-Holland en Zeeland, waaronder 11 op 24 september in het Verdronken Land van Saeftinghe. Er was een ringvangst van een **Krekelzanger** *Locustella fluviatilis* op 11 september bij Ter Wispel, Friesland. Er werden niet minder dan drie **Veldrietzangers** *Acrocephalus agricola* geringd en wel op 12 september bij Ter Wispel, op 17 september in de AW-duinen en op 14 oktober in Meijendel. **Waterrietzangers** *A paludicola* werden gevangen op 5 september in het Verdronken Land van Saeftinghe en op 11 september op Kampereiland, Overijssel; er waren late veldwaarnemingen op 25 september op Texel en op 3 en 4 oktober op de Maasvlakte. Een **'kleine spotvogel'** *A caligatus/rama* werd op 13 september kortstondig gezien in Dishoek, Zeeland. Een **Kleine Spotvogel** *A caligatus* liet zich van 9 tot 11 oktober zeer fraai bekijken op de Maasvlakte. Tot 10 oktober werd nog een 10-tal **Sperwergrasmussen** *Sylvia nisoria* doorgegeven waarvan vier veldwaarnemingen. De **Grauwe Fitis** *Phylloscopus trochiloides* van Koarnwertersân (Kornwerderzand), Friesland, was daar nog op 1 september. **Pallas' Boszangers** *P proregulus* verschenen op 7, 9 en 16 oktober op Texel, op 9 oktober op Vlieland, op 21 oktober op Schiermonnikoog (vangst) en in de Eemshaven, van 26 tot 28 oktober (ten minste twee) bij Petten, Noord-Holland, op 28 oktober bij Castricum, op 29 oktober bij Bloemendaal aan Zee en op 30 oktober op de Maasvlakte. Na de eerste **Bladkoningen** *P inornatus* op 9 september bij Katwijk aan Zee en 18 september op Terschelling volgden er van deze inmiddels gewone najaarsgast nog vele 10-tallen waarnemingen en vangsten in de rest van de periode. Op de ringbaan in Castricum werden er maar liefst 30 gevangen. Er waren **Raddes Boszangers** *P schwarzi* op 13 oktober bij Bloemendaal (vangst), op 19 oktober bij Castricum (vangst) en op 29 oktober bij Hargen aan Zee, Noord-Holland. **Bruine Boszangers** *P fuscatus* verbleven op 18 oktober bij de zuidpunt van de Brouwersdam, Zeeland, en op 24 en 25 oktober bij Lange Paal op Vlieland. **Siberische Tjiftjaffen** *P collybita tristis* werden gemeld op 12 oktober op Texel, op 24 oktober bij Paesens en op 28 oktober op de Eemshaven. **Kleine Vliegenvangers** *Ficedula parva* werden ontdekt op 29 september, 1 oktober en 14 en 15 oktober op Texel, op 30 september in Vlissingen, op 7 en 8 oktober op Vlieland, en op 10 oktober bij Stellendam, en een exemplaar vloog op 25 oktober langs telpost de Nolledijk bij Vlissingen. **Taigaboomkruipers** *Certhia familiaris* werden gevonden op 24 september en op 21 en 22 oktober op Vlieland, op 27 september bij Driene, Overijssel, op 2 en 20 oktober op Schiermonnikoog, op 13 oktober bij Nieuwolda, Groningen, op 14 oktober in de AW-duinen, en op 22 oktober in een mistnet in Helpman in de stad Groningen, Groningen (de soms lastige determinatie gaf hier en daar aanleiding om vraagtekens te plaatsen). Een Deense vogelaar ontdekte op 25 september een adult mannetje **Daurische Klauwier** *Lanius isabellinus* bij Den Hoorn op Texel. Deze was de volgende dag ook nog aanwezig. Eveneens op Texel maar dan op de noordpunt verbleef van 2 tot 9 september een eerstejaars **Roodkopklauwier** *L senator*. **Notenkrakers** *Nucifraga caryocatactes* werden gemeld op 17 september bij Bovensmilde, Drenthe, en in Amersfoort, Utrecht, op 20 september bij Roden, Drenthe, en op 21 september bij Ter Apel, Groningen. De discussie over de determinatie van een mannetje **Spreeuw** *Sturnus vulgaris* dat leek op een Zwarte Spreeuw *S unicolor* en vanaf 10 oktober in Oudeschild op Texel verbleef, hield de gehele periode aan. Vanaf 17 september werden c 11 juveniele **Roze Spreeuwen** *S roseus* waargenomen waarvan het merendeel op de Waddeneilanden. **Roodmussen** *Carpodacus erythrinus* doken op 13 september op bij IJmuiden, op 30 september op Texel en bij Bloemendaal (allebei vangsten), en op 15 oktober bij Westenschouwen, Zeeland. Er werden tot 15 oktober, maar voornamelijk in september, nog enkele 10-tallen **Ortolanen** *Emberiza hortulana* doorgegeven. De enige **Bosgors** *E rustica* van dit najaar vertoefde van 4 tot 7 oktober op camping Lange Paal op Vlieland. Vanaf 17 september werden 12 **Dwerggorzen** *E pusilla* genoteerd, veelal langstrekkend. Een mannetje **Zwartkopgors** *E melanocephala* verbleef van 16 tot 18 oktober in een groep Rietgorzen *E schoeniclus* aan de zuidpunt van de Brouwersdam en betrof het eerste najaarsgeval.

Ruud M van Dongen, Taalstraat 162, 5261 BJ Vught, Nederland
Remco Hofland, Aert van Neslaan 336, 2341 HN Oegstgeest, Nederland (remco.hofland @dutchbirding.nl)
Peter W W de Rouw, Schoolstraat 3-bis, 3581 PM Utrecht, Nederland

België

EENDEN TOT IBISSEN Van 15 tot ten minste 29 oktober verbleef een juveniele **Roodhalsgans** *Branta ruficollis* in de Uitkerkse Polders, West-Vlaanderen. Op zes plaatsen werden **Casarca's** *Tadorna ferruginea* opgemerkt, waaronder een groep van zes in het Schulensbroek, Limburg, op 6 september. De hele periode verbleef een

mannetje **Witoogeend** *Aythya nyroca* op De Gavers in Harelbeke, West-Vlaanderen, en op ten minste 29 oktober zwom hier een tweede. Andere mannetjes waren aanwezig van 3 september tot 5 oktober op de bezinkingsputten van Longchamps, Namur, van 10 tot 15 september bij Harchies, Hainaut, en van 15 september tot 2 oktober bij Sint-Lenaerts, Antwerpen. Op 14 oktober verschenen weer een mannetje en een vrouwtje in de Beneden-Netevallei in Lier-Duffel, Antwerpen; het vrouwtje bleef de rest van de maand. Op 3 oktober trok de eerste **Parelduiker** *Gavia arctica* langs De Panne, West-Vlaanderen; tussen 21 en 31 oktober werden langs de kust 19 exemplaren geteld. De enige **Roodhalsfuten** *Podiceps grisegena* werden gezien bij Oostende, West-Vlaanderen, op 4 september (langstrekkend), in Rijkevorsel, Antwerpen, op 11 september en in Hofstade, Vlaams-Brabant, op 1 en 19 oktober. Op 22 oktober zwommen twee **Kuifduikers** *P auritus* op De Kuifeend in de Antwerpse haven, Antwerpen. Een **Kuhls Pijlstormvogel** *Calonectris diomedea* vloog op 7 oktober langs De Panne. Langs Oostende werden **Vale Pijlstormvogels** *P mauretanicus* gemeld op 4 september en 29 oktober. In totaal werden 160 **Grauwe Pijlstormvogels** *P griseus* (met maximaal 68 langs De Panne op 3 oktober), slechts 23 **Noordse Pijlstormvogels** *P puffinus* en 10 **Vale Stormvogeltjes** *Oceanodroma leucorhoa* geteld. Het ontbreken van gunstige wind was daar uiteraard niet vreemd aan. Van 2 september tot ten minste 1 oktober verbleef een juveniele **Kuifaalscholver** *Phalacrocorax aristotelis* op de Kristallijnvijver in Lommel, Limburg, en op 10 september zwom er één op zee ter hoogte van Sint-Idesbald, West-Vlaanderen. Solitaire **Kwakken** *Nycticorax nycticorax* werden waargenomen in Hove, Antwerpen, op 15 september; in Lier-Duffel op 16 september; in Berlare-Zele, Oost-Vlaanderen, op 17 september; bij Pommeroeul, Hainaut, op 18 september; en in Kuurne, West-Vlaanderen, op 12 oktober. Van 5 tot 15 september vlogen bijna dagelijks maximaal zes **Koereigers** *Bubulcus ibis* over de Oesterputbrug bij Boekhoute, Oost-Vlaanderen, en van 19 tot 26 oktober passeerde hier vrijwel dagelijks nog één. Vanaf 9 september werden in het Blankaartbekken bij Woumen, West-Vlaanderen, geregeld maximaal zes foeragerende vogels opgemerkt. In Harchies, waar men van 1 tot 9 september dagelijks één vogel zag, verscheen op 7 september een groepje van zeven. Op 7 september liep er bovendien één in de Uitkerkse Polder, West-Vlaanderen, en op 11 september werd vermoedelijk hetzelfde exemplaar gezien bij Lissewege, West-Vlaanderen. De grootste concentraties **Kleine Zilverreigers** *Egretta garzetta* situeerden zich in Lissewege (maximaal 85 op 1 september); Het Zwin in Knokke, West-Vlaanderen (maximaal 71 op 8 september); Harchies-Hensies (28 op 4 september); de Achterhaven van Zeebrugge, West-Vlaanderen (maximaal 21 op 26 september); en de IJzermonding in Nieuwpoort, West-Vlaanderen (maximaal 10 op 28 september). **Grote Zilverreigers** *Casmerodius albus* werden 'schandalig veel' opgemerkt met grote(re) concentraties tot zelfs topaantallen in Harelbeke (11 op 12 oktober); Grimbergen, Vlaams-Brabant (15 op 29 oktober); Harchies (29 op 24 september); Roly, Namur (maximaal 32 op 14 oktober); Zonhoven, Limburg (maximaal 32 op 12 oktober); Wijnegem, Antwerpen (35 op 27 oktober); Zolder, Limburg (40 op 21 oktober); en Virelles, Hainaut (68 op 30 oktober). Tussen 8 en 25 september passeerden nog 30 **Purperreigers** *Ardea purpurea* met als dagmaximum een groep van zes over Bornem, Antwerpen, op 23 september. Op 1 oktober vloog er één over Bellem. In Zoutleeuw, Vlaams-Brabant, pleisterde een juveniele vogel van 16 september tot 15 oktober. Op 15 oktober vloog er nog één over Sint-Denijs, West-Vlaanderen, en vanaf 30 oktober (tot in november!) was er één aanwezig in de Viconiakleiputten in Stuivekenskerke, West-Vlaanderen. Tot 25 september was er gespreide doortrek van **Zwarte Ooievaars** *Ciconia nigra*: er waren 30 waarnemingen met 37 exemplaren. De laatste trokken op 8 oktober over Xhoris, Liège, en zelfs nog op 13 oktober over Munte, Oost-Vlaanderen. Op het totaal van **Ooievaars** *C ciconia* valt al moeilijker een cijfer te kleven maar het waren er vele 100en. De grootste groepen werden gezien in Klein Vorst, Antwerpen/Lommel (56 op 6 september), in Munsterbilzen, Limburg (50 op 2 september) en bij Froidchapelle, Hainaut (50 op 7 september). Van 1 tot 5 september verbleef een juveniele **Zwarte Ibis** *Plegadis falcinellus* in de Gentbrugse Meersen, Oost-Vlaanderen, en op 10 september werd dezelfde gezien bij Deinze, Oost-Vlaanderen.

583 Vorkstaartmeeuw / Sabine's Gull *Larus sabini*, adult, Oostende, West-Vlaanderen, 7 oktober 2006 *(Roland François)*

584 Noordse Boszanger / Arctic Warbler *Phylloscopus borealis*, Heist, West-Vlaanderen, 25 september 2006 *(Koen Verbanck)*

WOUWEN TOT SPECHTEN Er werden zeven **Zwarte Wouwen** *Milvus migrans* waargenomen waarvan vier in september. Op 10 september schoten **Rode Wouwen** *M milvus* echt goed uit de startblokken: in totaal volgden 162 met een bijzonder maximum van 22 in Hemroulle, Luxembourg, op 21 oktober. In Torgny werd op 8 oktober een overvliegende **Vale Gier** *Gyps fulvus* opgemerkt. Het was wel even slikken toen op 1 oktober een **Roodkopgier** *Cathartes aura* kwam aanvliegen op de telpost van de Bosdelkouter in Outgaarden, Vlaams-Brabant. De vogel was voorzien van de nodige toeters en bellen. Een **Slangenarend** *Circaetus gallicus* verscheen in het Mechels Broek bij Mechelen, Antwerpen, op 4, 6 en 8 september. Op 10 september trok er één over Lommel en op 1 oktober volgde een laatste waarneming op de Botrange, Liège. Een adult mannetje **Steppekiekendief** *Circus macrouros* vloog op 16 september over Sint-Denijs (op 20 augustus verbleef bij Hechtel, Limburg, waarschijnlijk een vrouwtje waarvan de identiteit nog wordt onderzocht). In de eerste helft van september werden nog negen **Grauwe Kiekendieven** *C pygargus* opgemerkt. Overtrekkende of langsvliegende **Ruigpootbuizerds** *Buteo lagopus* doken op in Harelbeke op 15 september; Eksaarde, Oost-Vlaanderen, op 21 september en 27 oktober; Grimbergen op 10 oktober; en Wellen, Limburg, op 29 oktober. Op 11 oktober was er een melding van een overvliegende lichte vorm **Dwergarend** *Aquila pennata* in Bree, Limburg. In totaal werden naar schatting 260 **Visarenden** *Pandion haliaetus* vastgesteld maar net zoals bij sommige andere soorten zijn dubbeltellingen niet uit te sluiten. **Roodpootvalken** *Falco vespertinus* vlogen op 12 september over Outgaarden en Jodoigne, Brabant-Walon, en op 15 september over Schulen. Een juveniel **Kleinst Waterhoen** *Porzana pusilla* liet zich van 29 september tot 5 oktober regelmatig vanaf grote afstand bekijken in de moerassen van Harchies. Waarnemingen van **Kwartelkoningen** *Crex crex* kwamen op 3 september van Elen, Limburg, en op 28 september van Maasmechelen, Limburg. Op 28 september was er een vondst aan de voet van de Sint-Baafskathedraal in hartje Gent, Oost-Vlaanderen, waar een paar Slechtvalken *F peregrinus* haar prooiresten dropt. Op 17 september werden in Wallonië de eerste 60 **Kraanvogels** *Grus grus* opgemerkt. Tussen 4 en 7 oktober waren enkele losse waarnemingen en de eerste echte trekpiek van enkele 1000en tekende zich af op 16 en 17 oktober met enkele nakomende groepen tot 26 oktober. De hoogste telling was op 16 oktober in Battice, Liège, met 500. In Vlaanderen werden slechts enkele kleine groepjes gezien en dan vrijwel uitsluitend in Limburg. Een **Griel** *Burhinus oedicnemus* liep op 5 september op de Scheldeschorren bij Kallo, Oost-Vlaanderen. Verrassend was te horen dat een **Herdersplevier** *Charadrius pecuarius* op 23 september doodleuk foerageerde in de Zilteweiden bij Oudenburg, West-Vlaanderen; naar verluidt droeg de vogel een gele kleurring. Overtrekkende **Morinelplevieren** *C morinellus* werden gezien in Genk op 2 september; in Brecht, Antwerpen, op 10 september; en in Zeebrugge op 21 september. Op 29 oktober pleisterde een laat exemplaar tussen de Goudplevieren *Pluvialis apricaria*

in De Moeren bij Adinkerke, West-Vlaanderen. Op 22 oktober werd in De Moeren bij Veurne, West-Vlaanderen, een **Aziatische Goudplevier** *P fulva* geringd. Een juveniele **vorkstaartplevier** *Glareola* pleisterde op 23 september in een groep Kieviten *Vanellus vanellus* in Gaurain, Hainaut, maar te kort om specifieke determinatie toe te laten… Juveniele **Gestreepte Strandlopers** *Calidris melanotos* verbleven van 2 tot 7 september op de Kristallijnvijver bij Lommel en van 2 tot 18 september op de bezinkingsputten van Longchamps. Op 12 oktober was er kortstondig één te zien bij Kieldrecht, Oost-Vlaanderen. Op 1 september vloog een **Grote Grijze Snip** *Limnodromus scolopaceus* over Hechtel-Eksel. **Grauwe Franjepoten** *Phalaropus lobatus* verschenen in Knokke op 2 september; Kieldrecht van 9 tot 12 september; Stuivekenskerke op 10 september; en de Uitkerkse Polders van 12 tot 14 september. De enige **Kleinste Jager** *Stercorarius longicaudus* voor de periode was een juveniele op 17 september langs de Duinbrug in Bredene, West-Vlaanderen. Op 7 oktober vlogen **Vorkstaartmeeuwen** *Larus sabini* langs Oostende en Koksijde, West-Vlaanderen, en vier langs De Panne. Op 27 oktober pleisterde kortstondig een juveniele in de Voorhaven van Zeebrugge en op 29 oktober trokken er twee langs De Panne. 'Jean', de adulte **Grote Burgemeester** *L hyperboreus* van de haven van Oostende, was op 12 september voor de 10e opeenvolgende winter terug en bleef de hele periode. In De Gavers in Harelbeke werden op 9 september twee langsvliegende **Reuzensterns** *Sterna caspia* waargenomen: een adulte en een eerste-winter. Op 2 september werd gezien hoe de groep van 20 **Bijeneters** *Merops apiaster* van het Provinciaal Domein Puyenbroek in Wachtebeke, Oost-Vlaanderen, hoogte won en in zuidoostelijke richting verdween. Een groep van 15 die op 5 september over Merelbeke, Oost-Vlaanderen, trok, had wellicht betrekking op dezelfde vogels. **Middelste Bonte Spechten** *Dendrocopos medius* werden gezien in Geistingen, Limburg; Maasmechelen; en Oostmalle, Antwerpen.

LEEUWERIKEN TOT GORZEN Overtrekkende **Kortteenleeuweriken** *Calandrella brachydactyla* werden herkend in Lier op 14 oktober en in Wingene op 25 oktober. Er volgde een ringvangst in Zele op 30 oktober. Verrassend was de waarneming van twee **Roodstuitzwaluwen** *Cecropis daurica* die in de vooravond van 10 oktober over de Achterhaven van Zeebrugge vlogen. Vanaf 26 september was er een gespreide doortrek van **Grote Piepers** *Anthus richardi*: er werden 23 waarnemingen ontvangen. Met 60 septemberwaarnemingen deed **Duinpieper** *A campestris* het minder goed dan vorig jaar. In oktober volgden nog waarnemingen in Brecht (twee op 1 oktober), Lier (9 oktober) en Kapellen (26 oktober). Tussen 23 september en 17 oktober werden 23 **Roodkeelpiepers** *A cervinus* opgemerkt en op 30 oktober volgde een laatste waarneming in Zeebrugge. De derde **Blauwstaart** *Tarsiger cyanurus* voor België betrof een ringvangst in Korbeek-Lo, Vlaams-Brabant, op 17 oktober. Op 27 oktober werd een mannetje **Aziatische Roodborsttapuit** *Saxicola maurus* gezien en gefotografeerd in Hemptinne, Namur. Op heel wat plaatsen werden weer **Cetti's Zangers** *Cettia cetti* gehoord en/of gezien. Datzelfde geldt ook voor **Graszangers** *Cisticola juncidis*, waarvan er op 9 september 22 werden geteld in de Achterhaven van Zeebrugge! Er waren veldwaarnemingen van **Waterrietzangers** *Acrocephalus paludicola* in het Bos van Aa in Zemst, Vlaams-Brabant, op 3 september en in de Achterhaven van Zeebrugge op 11 september. De laatste waarneming was een ringvangst in Zulte, West-Vlaanderen, op 17 september. Leuk was de ringvangst van een **Veldrietzanger** *A agricola* bij Genk op 22 september. Zowel op 1 als 17 september werd een **Grote Karekiet** *A arundinaceus* gevangen, een soort die nu duidelijk lijkt te verdwijnen. Eerste-winter **Sperwergrasmussen** *Sylvia nisoria* werden gezien in de Voorhaven van Zeebrugge op 17 september en op 5 oktober. Klassiek bij deze soort is het overwicht aan ringvangsten: in Hamme, Oost-Vlaanderen, op 1 september; bij Veurne op 2 september; in Oostduinkerke, West-Vlaanderen, op 16 september; en in Awirs, Liège, op 19 oktober. Een **Noordse Boszanger** *Phylloscopus borealis* die van 24 tot 26 september in het Haagje (Sashul) in Heist verbleef liet zich vaak erg goed bekijken. Er verschenen **Pallas' Boszangers** *P proregulus* in Blankenberge op 15 oktober; in Zeebrugge op 28 oktober; en in Koksijde op 30 oktober. Bovendien waren er in het binnenland twee ringvangsten: in Nerem, Limburg, op 22 oktober en in Boutersem, Vlaams-Brabant, op 25 oktober. Voor **Bladkoningen** *P inornatus* betekende 2006, na het recordjaar 2005, het op twee na beste jaar ooit. De eerste verschenen op 24 september en gespreid over de periode werden er c 89 gemeld. Hiervan kwam het overwicht (c 41) van de drie oostelijke kustgemeenten (de Oostkust) maar de 24 waarnemingen uit het binnenland zijn niet te verwaarlozen. Een **Bruine Boszanger** *P fuscatus* toonde zich op 28 oktober wat moeizaam in de Baai van Heist. Tjiftjaffen *P collybita* met de 'klassieke' kenmerken van **Siberische Tjiftjaf** *P c tristis* werden gezien of geringd in Heist op 26 oktober; Genk op 28 oktober; en Kalmthout, Antwerpen, op 30 oktober. Tegen het klassieke patroon in werden **Kleine Vliegenvangers** *Ficedula parva* alleen opgemerkt in het binnenland: op 10 september in Kapelle-op-den-Bos, Vlaams-Brabant, en op 25 september in Merelbeke. Een **Taigaboomkruiper** *Certhia familiaris* werd op 14 oktober geringd in Tongeren en op 26 en 27 oktober werd een exemplaar gezien in het Visserskruis in Zeebrugge. Er werden in totaal meer dan 103 **Buidelmezen** *Remiz pendulinus* waargenomen met een duidelijk oostelijk zwaartepunt: als grootste concentratie verbleven er 26 op de bezinkingsputten bij Hollogne-sur-Geer, Liège, op 7 oktober. Bij Amay, Liège, werd geregelde doortrek vastgesteld met 16 op 10 oktober en 15 andere op 22 oktober. Tussen 1 en 30 september werden 11 **Grauwe Klauwieren** *Lanius collurio* gezien. Op 26 oktober was een late aanwezig in Heist. Vanaf 22 september verschenen her en der weer de nodige **Klapeksters** *L excubitor*: er waren 25 meldingen en in navolging van vorig jaar hadden intensieve trektellingen met 14 meldingen ook dit jaar weer een belangrijke inbreng. Op 24 september

verbleef kortstondig een juveniele **Roze Spreeuw** *Sturnus roseus* bij de Kleiputten van Heist. Tussen 2 september en 8 oktober werden 19 **Ortolanen** *Emberiza hortulana* waargenomen, merendeels overtrekkende vogels. Een **Dwerggors** *E pusilla* trok op 29 oktober laag en luid roepend over de telpost in De Maten in Genk.

De hulp van al diegenen die (hun) waarnemingen inspraken op de Natuurpunt-Vogellijn was hier onontbeerlijk. De Natuurpunt-Vogellijn is alleen vanuit België bereikbaar op het nummer 0900-00194 (EUR 0.45/min), de Natuurpunt-Inspreeklijn is te bereiken op 0800-11194 (gratis). De Waalse gegevens werden in hoofdzaak geput uit de AVES-website.

Gerald Driessens, Pastoriestraat 16, 2500 Lier, België (gerald.driessens@natuurpunt.be)

DB Actueel

Afwijkende spreeuw op Texel houdt gemoederen bezig
Op dinsdag 10 oktober 2006 maakte Wim Wiegant omstreeks 16:00 melding van een mogelijke Zwarte Spreeuw *Sturnus unicolor*. Hij had de vogel ontdekt tijdens een bezoek aan de haven van Oudeschild op Texel, Noord-Holland, waar hij en zijn partner juist de snackbar hadden bezocht. Omdat de traditionele Dutch Birding-vogelweek op het eiland in volle gang was, arriveerden in korte tijd andere vogelaars, die zich mengden in de discussie over de determinatie en zich toelegden op de documentatie. De meesten waren het erover eens dat het een opvallend donkere spreeuw *Sturnus* betrof die sterk deed denken aan Zwarte Spreeuw. Ongeveer twee uren later verdween hij uit beeld, maar aan de hand van de gemaakte foto's konden velen zich die avond verder buigen over de determinatie.

De foto's toonden onder meer een gele snavel met een blauwgrijze basis, een kenmerk van adulte mannetjes spreeuwen in het voorjaar, evenals een donkere iris, voor alle seizoenen een kenmerk van mannetjes. Ook was te zien dat het verenkleed overwegend zwart was, met slechts onopvallende lichte 'spikkels en randjes' op delen van het verenkleed, afwijkend van alle aanwezige Spreeuwen *S vulgaris* die hun bont gevlekte verse kleed droegen. Daar kwam bij dat een groot deel van het verenkleed een blauwpaarse glans vertoonde en dat de poten opvallend roze waren met witachtige nagels. Het feit dat de vogel zijn volledige rui nog niet had afgerond was onder meer te zien aan de nog groeiende p7 (handpennen van binnen naar buiten genummerd) en de twee gebleekte en gesleten oude buitenste lange handpennen (p8 en p9) in beide vleugels (p10 is bij spreeu-

585 Spreeuw / Common Starling *Sturnus vulgaris*, mannetje, Oudeschild, Texel, Noord-Holland, 11 oktober 2006 *(René Pop)*

wen gereduceerd, nog korter dan de handpendekveren, en is in het veld niet of nauwelijks zichtbaar).

Voor sommigen was de conclusie na het zien van deze beelden helder; de Oudeschild-spreeuw kon geen adult mannetje Zwarte Spreeuw zijn, omdat die volgens de meeste bronnen ongeacht de tijd van het jaar geen wit op de veerranden vertoont. Bovendien was van de kenmerkende verlengde veren op de kop, hals en borst van adulte mannetjes Zwarte Spreeuw geen spoor terug te vinden. Wat structuur betreft week de Oudeschild-spreeuw niet af van de andere aanwezige Spreeuwen. Anderen zagen in dit gegeven echter nog geen aanleiding om Zwarte Spreeuw als kandidaat af te schrijven omdat er geen verklaring was voor het afwijkende uiterlijk.

De volgende dag werd de 'zwarte spreeuw' weer in de haven van Oudeschild aangetroffen. Aanvullende kenmerken konden worden genoteerd, zoals het gedrag. Hij zat vaak op daken van gebouwen, joeg soms Spreeuwen weg, foerageerde op de grond tussen mensen en zong met enige regelmaat. Het leek wel alsof hij de openingstijden van een lokale patatzaak kende en ook bij het binnenlopen van een garnalenboot in de haven was hij er als de kippen bij. Dit alles bracht de determinatie echter niet veel verder. Veel waarnemers zagen wel donkere ondervleugeldekveren, iets wat sterk zou pleiten voor Zwarte Spreeuw, maar volgens anderen was hierin geen verschil te zien met gewone Spreeuw.

Allerlei informatiebronnen, waaronder enkele buitenlandse experts, werden in de daarop volgende dagen geraadpleegd, waardoor een aantal interessante punten aan het licht kwam. Er werd studie gemaakt van de enigszins gelijkende 'Scilly-spreeuw', die vanaf begin 1998 enige tijd de gemoederen bezighield op St Mary's, Scilly, Engeland (cf Birding World 11: 53-57, 1998), en de uiterlijke kenmerken van oostelijke ondersoorten van Spreeuw en van hybriden Spreeuw x Zwarte Spreeuw werden besproken. Maar dit alles leverde geen bewijzen op die de determinatie als Zwarte Spreeuw rechtvaardigden. De lichte tekening op lichaams- en vleugelveren en het ontbreken van verlengde kop-, hals- en borstveren bleken telkens struikelblokken, evenals het feit dat de snavel nog geel was in oktober. De geluiden die hij voortbracht waren evenmin reden om aan Zwarte Spreeuw te denken. Noch het opvallende *tepiet*-roepje, noch de zang, waarin imitaties van Texelse soorten waren te horen, pleitten voor Zwarte Spreeuw.

De verklaring voor het afwijkende uiterlijk van de Oudeschild-spreeuw moet waarschijnlijk worden gezocht in een hormonale afwijking. De vogel zou bijvoorbeeld door een gezwel op de testis een verkeerde hormoonspiegel kunnen hebben (Kees Roselaar in litt). Hierdoor vertoont hij resten van het uiterlijk in de broedtijd, zoals de kleur van de naakte delen, en wordt wellicht de late rui verklaard. De weinige lichaamsveren met witte randjes bleken zo veel sleet te vertonen dat ze niet vers geruid konden zijn en konden dus daarom al onmogelijk betrekking hebben op een mannetje Zwarte Spreeuw. Na weken van studeren deelden veel vogelaars de visie dat het een afwijkende Spreeuw met verlate rui en sterk gesleten witte veertoppen moest zijn. Maar er waren er ook die de hoop op een Zwarte Spreeuw (of ten minste een hybride) nog niet hadden opgegeven, daarbij gesteund door e-mails van twee Spaanse vogelaars. Feit is wel, dat het fraaie statief dat Swarovski in het kader van de Dutch Birding-vogelweek had uitgeloofd voor de 'Soort van de week' aan Wims neus voorbijging. Eigenlijk had hij dat statief sowieso verdiend voor het feit dat dit het 25e jaar was waarin hij een Dutch Birding-vogelweek bezocht. Ten noorden van Frankrijk is maar één geval van Zwarte Spreeuw aanvaard, van een volwassen mannetje dat van 18 april tot 27 mei 2002 en van 15 april tot 18 mei 2003 werd gezien en gefotografeerd in Denemarken.
ROY SLATERUS

ABERRANT STARLING From 10 October 2006, a starling *Sturnus* at Oudeschild, Texel, Noord-Holland, the Netherlands, provoked a lot of discussion and drew many birders to the island. It was a black starling with spring-like yellow bill with bluish base and bright pink legs. Because of the colour of bill base and dark iris it could be identified as a male older than one year. Small white feather markings were only visible on the wing-coverts, underparts and undertail-coverts. Most feathers, including those with white fringes on the underparts, were (very) worn, while the wing showed active primary moult, with p7 (numbered from the inside) still growing. From a distance, the bird looked all-black and much like a Spotless Starling *S unicolor*. However, after the annual complete moult, Spotless and Common Starling *S vulgaris* should have a fresh plumage at this time of year with darkish bill and legs. Adult male Spotless cannot have any white markings on worn body feathers (very small pale markings, if any, should wear off quickly). These anomalies indicated that it was an aberrant Common Starling *S vulgaris* probably with some kind of hormonal problem causing bright colours of bare parts and a delayed moult.

Vale Gierzwaluw op Vlieland: langverwachte nieuwkomer voor Nederland Vrijdag 20 oktober 2006 was de aftrap van het derde en laatste 'Deception Tours' weekend op Vlieland, Friesland. De twee eerdere weekenden waren rustig verlopen maar dit maakte de gebruikelijke speculaties en wilde fantasieën onder de c 15 deelnemers op de veerboot naar het eiland alleen maar groter…

Een paar uren later bevond ik (Nils van Duivendijk) me in de duinen tussen Duinkersoord en Lange Paal. Rond 14:00 werd mijn aandacht getrokken door een op grote afstand vliegende kleine vogel boven de zeereep. Door de kijker bleek het een gierzwaluw *Apus*! De vlucht leek wat traag en de bovenzijde wat bruinachtig: Alpengierzwaluw *A melba*?!? Maar erg groot leek hij niet. Net toen ik mijn telescoop wilde opzetten verdween hij achter de laatste duinenrij richting strand. Ik belde direct Diederik Kok van wie ik wist dat hij in de buurt was. In een lange sprint fietste ik naar de strandopgang van Lange Paal met het bange vermoe-

586 Vale Gierzwaluw / Pallid Swift *Apus pallidus*, Vlieland, Friesland, 20 oktober 2006 *(Bas van den Boogaard)*

587 Vale Gierzwaluw / Pallid Swift *Apus pallidus*, Vlieland, Friesland, 21 oktober 2006 *(Bas van den Boogaard)*

den de vogel nooit meer terug te zien – maar staand op het hoogste duin vond ik hem direct boven de zeereep ter hoogte van de Nieuwe Kooi. Door de telescoop viel nu een wat breed overkomende handvleugel op; door het donkere weer was de kleur niet te zien. Gelukkig zag ik Diederik over het fietspad naderen. Het bleek nu zelfs om twee gierzwaluwen te gaan! Met z'n tweeën pikten we één vogel op die iets dichterbij kwam. We volgden deze enige tijd door de telescoop totdat Diederik zei 'hij is bruin!'. 'Ja wat, zwartbruin als een Gierzwaluw of anders?' riep ik. 'Nee, lichter bruin...'. Juist, dat dacht ik dus ook, maar ik durfde niet aan Vale Gierzwaluw *A pallidus* te denken (zou een juveniele Gierzwaluw *A apus* door enige bleking zo laat in het jaar er ook niet zo uit kunnen zien?). We waarschuwden snel telefonisch wat mensen. We volgden nu de tweede vogel en de adrenalinespiegel zakte en de hartslag normaliseerde: we zagen nu niks wat niet past op gewone Gierzwaluw en twee verschillende soorten leek niet realistisch... Beide vogels vlogen eerst een paar kilometers naar het oosten en daarna een stuk de zee op maar steeds als we ze voorgoed dachten kwijt te raken, kwamen ze weer terug. Eindelijk kwam een vogel wat dichterbij, en nu waren we er vrij zeker van dat dit echt een gewone Gierzwaluw was.

Net toen we er zo'n beetje klaar mee waren, kwamen de vogels wéér dichterbij – toch nog even kijken. De bij elkaar vliegende vogels bekijkend vielen ons toch echt verschillen op: één had meer contrast op de boven- en ondervleugel en een iets andere jizz met stompere vleugelpunten en leek af en toe een bruine tint te hebben. Onze gedachten aan Vale Gierzwaluw namen weer toe – of liepen we onszelf nu gek te maken? Lichtelijk in paniek probeerden we tegelijkertijd beide vogels niet kwijt te raken, kenmerken vast te stellen, foto's te maken en anderen te bellen. Jeroen de Bruyn zag beide vogels vervolgens vanaf Duinkersoord en merkte ook de verschillen op. Inmiddels hadden wij versterking gekregen van de eerste gearriveerde vogelaars, waaronder Han Zevenhuizen die op het strand enkele plaatjes wist te maken. Het schermpje van de camera toonde nog meer kenmerken die pleiten voor Vale – het werd nu echt serieus! De pieperboodschap 'mogelijk' werd geplaatst. Juist toen ik aan het piepen was, vloog de vogel voor het eerst dichtbij langs en Diederik zag hem zodanig dat hij nu zeker was van determinatie als Vale. Ongeloof overheerste maar we stonden echt naar een nieuwe soort voor Nederland te kijken!

Terwijl de rappe fotografen Bas van den Boogaard en Han hun waarde bewezen, wisten alle Vlielandgangers de vogel te zien. Hij bleef zich steeds verplaatsen langs de kustlijn, soms vrij ver de zee opvliegend en vaak in direct gezelschap van de Gierzwaluw. Er trad nog even verwarring op toen zich een derde gierzwaluw aandiende maar deze werd snel ontmaskerd als ook een gewone Gierzwaluw. Op afstand bleek regelmatig hoe moeilijk het was de juiste vogel eruit te pikken – zeker met het grijze weer was de bruintint maar af en toe te zien. Kenmerken die beter bruikbaar bleken waren de structuur, het contrast in de vleugels, het donkere 'zadel' op de bovendelen en de grote lichte keelvlek die het voorhoofd bereikt. Terwijl Jeroen, Hans ter Haar en ik nog op de duintop zaten kwam hij onder ideale omstandigheden dichtbij langsvliegen. Nu zagen we ook de laatste nog ontbrekende details: lichte kop met duidelijk masker, lichtere stuit met duidelijke schubben en contrasterend met de donkerdere bovendelen, licht ruitpatroon op de buik en de koffie-met-melk-kleur. Het laatste stukje ongeloof was nu ook bij mij weg! De Vale Gierzwaluw werd rond 16:00 voor het laatst die dag met zekerheid gezien. Dezelfde avond staken nog c 10 vogelaars over van de vaste wal om de volgende ochtend met het eerste licht op de plek te kunnen zijn.

De volgende dag maakten c 35 vogelaars de oversteek met de boot van 09:00 vanuit Harlingen, Friesland. De nodige consternatie ontstond aan boord als rond 10:05 hoogstwaarschijnlijk de Vale Gierzwaluw vanaf de veerboot in het zeegat tussen Terschelling en Vlieland wordt gezien – net te ver en te kort om hele-

maal zeker te kunnen zijn. Rond 11:00 zagen de al op het eiland aanwezige waarnemers de vogel weer boven de duinenrij vliegen (waarschijnlijk had hij een grote lus gemaakt over de oostkant van het eiland). Om 12:00 konden ook de mensen die van de boot kwamen opgelucht ademhalen als eerst de twee Gierzwaluwen en daarna de Vale Gierzwaluw langzaam van west naar oost over de duinenrij vlogen en op zee uit beeld verdwenen. Gedurende de rest van de dag werd de Vale met (soms lange) tussenposen langs de Noordzeekust gezien, veelal in de omgeving van Hotel Seeduyn. De c 10 mensen die met de middagboot arriveerden zagen hem vanaf c 18:00 in het laatste licht geruime tijd boven het strand heen en weer vliegen. Pas om 18:40 was hij op korte afstand ten noorden van het hotel uit beeld verdwenen. De volgende dag werd hij, tot teleurstelling van c 35 overgekomen vogelaars, niet meer waargenomen; ook de beide Gierzwaluwen waren verdwenen.

Indien aanvaard betreft dit het eerste geval van Vale Gierzwaluw. De lastige veldherkenning en het feit dat het moeilijk is om deze soort met goede foto's te documenteren vormden bij vorige meldingen obstakels voor toetreding tot de Nederlandse lijst. Ons geluk was dat deze vogel 'gevangen zat' op een eiland. Daardoor was hij niet alleen twitchbaar maar ook goed te bestuderen en te fotograferen. Het was opmerkelijk dat eind oktober meer late Gierzwaluwen dan gebruikelijk in Nederland werden opgemerkt. Ook volgden er nog enkele meldingen van mogelijke Vale Gierzwaluwen elders in het land. De meest serieuze betrof een exemplaar langs de Nolle bij Vlissingen, Zeeland, op 25 oktober. Nils van Duivendijk & Diederik Kok

Pallid Swift On 20-21 October 2006, a Pallid Swift *Apus pallidus* was present on Vlieland, Friesland, the Netherlands. It was seen by c 15 birders staying on the island on the first day and successfully twitched by c 60 birders on the second day, when it flew in large circles along the coast and over the sea. On the first day, it was often seen with one or two Common Swifts *Apus apus*, allowing direct comparison of shape, plumage and flight actions. If accepted, it will be the long-awaited first record of this species for the Dutch list. Several reports in previous years proved to be insufficiently documented but this bird was well photographed.

Eindelijk Rotszwaluwen Vanaf 20 oktober 2006 werden in Zweden binnen enkele dagen meerdere Rotszwaluwen *Ptyonoprogne rupestris* waargenomen: een groepje van vier, een groepje van drie en een enkeling. Er waren slechts twee eerdere gevallen. Op 22 oktober was er ook een melding in Engeland. Het zeer zachte weer dat de eerste helft van november 2006 kenmerkte deed de hoop stijgen dat ook Nederland eindelijk een aanvaardbare en twitchbare Rotszwaluw ten deel zou vallen. Na een lange aanloop zou dat inderdaad lukken... Het begon met een melding van twee exemplaren door Rob Baars en Sietske Sybrandi op het Steigereiland in IJburg, Amsterdam, Noord-Holland, in de late middag van zondag 5 november. Hun beschrijving paste uitstekend op Rotszwaluw; zelfs de witte vlekjes in de staart werden vastgesteld. De volgende ochtend hoopten c 20 vogelaars op een klein wonder maar dat gebeurde niet. Twee dagen later, op 7 november, zagen Bert Kleijn, Cor Oskam, Leen van Ree en Wout Vuyk bij Vogelringstation Nebularia in Westenschouwen, Zeeland, rond 13:00 gedurende c 2 min twee Rotszwaluwen aan de rand van een naaldbos voordat ze in noordelijke richting wegvlogen. Eén van beide werd goed door de verrekijker bekeken waarbij de witte staartvlekken werden gezien. Tevens werd een Huiszwaluw *Delichon urbicum*-achtige roep gehoord en bij deze waarneming laat de beschrijving eigenlijk geen ruimte voor twijfel over de determinatie. De volgende zet in het kat-en-muisspel tussen vogelend Nederland en deze soort kwam van Maarten Hotting die op dinsdag 14 november rond 09:30 vanuit zijn klaslokaal tijdens een les Nederlands in schoolgebouw De Titaan in Hoorn, Noord-Holland, kortstondig een forse bruine zwaluw zag vliegen. Hij dacht gelijk aan een Rotszwaluw en plaatste zijn melding op de website www.waarneming.nl. Het duurde tot c 15:00 voordat deze werd doorgegeven op het semafoonsysteem. Nick van der Ham en Sander Lagerveld gingen hun geluk beproeven en zagen tot hun verbazing kort nadat ze bij de school waren aangekomen om c 15:55 drie zwaluwen vliegen: een Boerenzwaluw *Hirundo rustica*, een vermoedelijke Rotszwaluw en een ongedetermineerde zwaluw. De vogels vlogen op c 500 m afstand bij de IJsselmeerdijk en konden niet goed worden bekeken; wel slaagde SL er in om enkele seconden aan videoopnamen te maken van de vermeende Rotszwaluw. De beelden boden misschien geen 100% zekerheid over de determinatie maar de aanwezigheid van minimaal één Rotszwaluw leek wel erg waarschijnlijk. De volgende ochtend beproefden c 30 vogelaars vanaf het eerste licht hun geluk bij de school en op de dijk maar deze dag werd geen enkele zwaluw gemeld, misschien mede doordat het vrij slecht weer was. Op 16 november zag MH om 15:45 opnieuw een Rotszwaluw vliegen vanuit zijn klaslokaal. De volgende dag trokken daarom wederom enkele 10-tallen vogelaars naar Hoorn en van hen werden c 20 volhouders (sommigen waren er al vanaf 08:00) en enkele gelukkige 'laatkomers' om 14:30 beloond met een fraaie vliegshow van 2-3 min van twee Rotszwaluwen rond het schoolgebouw. Nu konden ze eindelijk goed worden gedocumenteerd door foto's van Michel Veldt. Veel vogelaars snelden na de melding nog voor donker naar de plek maar ze kwamen voor niets – de zwaluwen werden niet meer gezien. Op zaterdagochtend 18 november kwamen 10-tallen vogelaars 's ochtends naar Hoorn. De meesten bleven angstvallig bij De Titaan maar anderen waren zo dapper of ongeduldig om in de omgeving te gaan zoeken. Dat leverde al snel succes op – rond 09:10 zagen Remco Hofland en Chris Quispel de twee Rotszwaluwen vliegen rond de flatgebouwen aan de dijk ten westen van de schouwburg. Hier kwamen ze enkele keren terug en lieten zich door talloze vogelaars fraai bekijken en op video en foto vastleggen. Rond 10:30 gingen ze zitten

588 Rotszwaluw / Eurasian Crag Martin *Ptyonoprogne rupestris*, Hoorn, Noord-Holland, 18 november 2006 *(Arnoud B van den Berg)*

589 Rotszwaluw / Eurasian Crag Martin *Ptyonoprogne rupestris*, Hoorn, Noord-Holland, 18 november 2006 *(René van Rossum)*

op een spuwer hoog op een flatgebouw aan De Weel. Enkele fotografen konden vanuit het appartement op de bovenste verdieping prachtige foto's maken. Rond 11:00 verplaatsten de twee zwaluwen zich naar het oude centrum waar ze rond de Grote Kerk vlogen. Rond 11:45 werden ze op hun oude plek bij De Titaan gezien – en zo ging het de rest van de dag (tot c 16:30) door, waarbij enkele 100-en vogelaars de soort eindelijk op hun Nederlandse lijst konden bijschrijven. Ook de volgende dagen werden de vogels nog gezien, ten minste tot en met 24 november.

De vogels van Hoorn zijn de eerste uitgebreid gedocumenteerde Rotszwaluwen voor Nederland. Het is afwachten of de waarnemingen van IJburg en Westenschouwen aanvaardbaar zijn voor de Commissie Dwaalgasten Nederlandse Avifauna (CDNA). Er lijkt sprake van een influx van zes vogels maar mogelijk hebben de waarnemingen van IJburg en Hoorn, beide aan de IJsselmeerkust en op c 30 km afstand, betrekking op dezelfde vogels. Eerdere waarnemingen in Nederland zijn nog in behandeling bij de CDNA of zijn afgewezen omdat de documentatie niet volledig was. Ze hadden alle betrekking op exemplaren die in de trektijd langsvlogen waardoor het, anders dan bij de vogels van Hoorn, vrijwel onmogelijk was om ze uitgebreid te bestuderen, beschrijven en/of fotograferen. Maarten Hotting & Enno B Ebels

Eurasian Crag Martins On 5 November 2006, two Eurasian Crag Martins *Ptyonoprogne rupestris* were seen briefly by one birder and his friend at IJburg, Amsterdam, Noord-Holland, the Netherlands. On 7 November, two were seen and heard calling by four observers at Westenschouwen, Zeeland. From 14 to at least 24 November, one to two were seen (almost) daily but, in the first days, briefly at Hoorn, Noord-Holland (c 30 km north of IJburg). On 17 November, the first photographs of the two birds at Hoorn were made and, on 18 November, the two were seen for long periods, both at rest and in flight, during the

590 Rotszwaluwen / Eurasian Crag Martins *Ptyonoprogne rupestris*, Hoorn, Noord-Holland, 18 november 2006 *(Chris van Rijswijk)*

whole day by several 100s of birders. The Hoorn birds are the first to be documented by photographs and video and may constitute the first record for the Netherlands, unless the reports in the first week of November (or reports from previous years) will be accepted.

Huisgierzwaluw te IJmuiden Op maandag 20 november 2006 om 13:00 zag ik (Bram Couperus) vanuit een raam op de begane grond van het gebouw Wageningen IMARES (voorheen bekend als het RIVO-gebouw) aan de Haringkade te IJmuiden, Noord-Holland, een 'gierzwaluw' langsvliegen. De vogel vloog in oostelijke richting op c 20-25 m hoogte vlakbij de rotonde aan de zuidwesthoek van de Vissershaven. Ik waarschuwde direct mijn collega, Joep de Leeuw, en met onze verrekijkers gingen we naar buiten. We zagen daar direct de vogel, die op 20 m hoogte boven de rotonde rondvloog en vervolgens landde op de betonnen wand van de vervallen oude watertoren die bij de rotonde staat. Bij het landen hield hij zijn vleugels naar achteren en zijn staart gespreid. De witte stuit viel daarbij direct op. Onmiddellijk steeg ons adrenalinepeil, maar beiden hadden we geen ervaring met een 'gierzwaluw met witte stuit'. Hij zat op een hoogte van c 20 m met zijn rug naar de Trawlerkade en c 25 cm onder een dichtgetimmerd raam. Vervolgens kroop hij omhoog tot iets onder een betonnen raamkozijn. Daarbij hield hij zijn lichaam verticaal, iets naar rechts hellend, en met de kop naar rechts gedraaid. Vanaf de overzijde van de weg, op een afstand van 30-40 m, konden we de vogel goed bekijken. Het verenkleed was donker grijsbruin en de zeer opvallende helder witte stuitvlek groot en bijna vierkant, aan de onderrand iets afgerond. De vleugelpunten staken tot ver voorbij de donkerbruine staart, die met de punt tegen de muur werd aangedrukt. De kop was rond met een zeer korte donkere snavel. Het voorhoofd en de wenkbrauwstreep waren iets lichter. Verder had de vogel een opvallende, grote, zeer lichte keelvlek. Zelfs met het blote oog waren de witte stuitvlek en de lichte keel te zien. Na 2-3 min vloog hij recht over ons heen in noordwestelijke richting over de haven weg. Hierbij waren het kleine formaat met kortere, rechtere vleugels dan Gierzwaluw *Apus apus* en de korte, recht afgesneden of hooguit licht ingekeepte staart goed te zien. Terug op de werkplek gingen we naarstig op zoek naar determinatiewerken. BC keek op internet en JdL vond een oude 'Heinzel' en een minuut later zeiden we tegelijk respectievelijk 'Little Swift' en 'Huisgierzwaluw' en meldde JdL de waarneming van Huisgierzwaluw *A affinis* mede namens BC direct om 13:23 op VogelnetZuidKennemerland. Later werd een gedetailleerde beschrijving gemaakt en bevestigden andere geraadpleegde veldgidsen en *Birds of the Western Palearctic* de determinatie. Het internetbericht bereikte in korte tijd ook het semafoonsysteem en de rest van de middag kwamen enkele 10-tallen vogelaars naar de Vissershaven. De vogel kon in regenachtige omstandigheden echter niet meer worden teruggevonden. Indien aanvaard betreft dit het eerste geval voor Nederland. Een melding van Terschelling, Friesland, op 17 mei 2001 (Dutch Birding 23: 237, plaat 268-269, 2001) werd niet door de Commissie Dwaalgasten Nederlandse Avifauna aanvaard omdat geen beschrijving werd ingediend. Bram Couperus & Joep de Leeuw

Little Swift On 20 November 2006, a Little Swift *Apus affinis* was seen for 5 min by two birders at IJmuiden, Noord-Holland, the Netherlands. The bird flew at a height of c 20 m over a road and landed on an old concrete tower. After a few minutes, it flew off again and disappeared over the harbour. If accepted, it will be the first for the Netherlands. A previous report from Terschelling, Friesland, on 17 May 2001 was not submitted and, therefore, not accepted.